

cse371/mat371
LOGIC

Professor Anita Wasilewska

Fall 2016

LECTURE 10a

Chapter 10
CLASSICAL AUTOMATED PROOF SYSTEMS

PART 3: GENTZEN SYSTEMS

Gentzen Sequent Calculus GL

The proof system **GL** for the classical propositional logic presented now is a **version** of the original **Gentzen** (1934) systems **LK**.

A **constructive** proof of the **Completeness Theorem** for the system **GL** is very similar to the proof of the Completeness Theorem for the system **RS**

Expressions of the system are like in the **original Gentzen** system **LK** are Gentzen **sequents**

Hence we use also a name **Gentzen sequent calculus** for it

Gentzen Sequent Calculus GL

Language of **GL**

$$\mathcal{L} = \mathcal{L}_{\{ \cup, \cap, \Rightarrow, \neg \}}$$

We **add** a new symbol to the alphabet \longrightarrow called a **Gentzen arrow**

The **sequents** are built out of **finite sequences** (empty included) of formulas, i.e. elements of \mathcal{F}^* , and the additional symbol \longrightarrow

We **denote**, as in the **RS** system, the finite sequences of formulas by Greek capital letters

$$\Gamma, \Delta, \Sigma, \dots$$

with indices if necessary

Gentzen Sequents

Definition Any expression

$$\Gamma \longrightarrow \Delta$$

where $\Gamma, \Delta \in \mathcal{F}^*$ is called a **sequent**

Intuitively, we **interpret semantically** a sequent

$$A_1, \dots, A_n \longrightarrow B_1, \dots, B_m$$

where $n, m \geq 1$, as a formula

$$(A_1 \cap \dots \cap A_n) \Rightarrow (B_1 \cup \dots \cup B_m)$$

Gentzen Sequents

The sequent

$$A_1, \dots, A_n \longrightarrow$$

(where $m \geq 1$) means that $A_1 \cap \dots \cap A_n$ yields a **contradiction**

The sequent

$$\longrightarrow B_1, \dots, B_m$$

(where $m \geq 1$) means semantically $T \Rightarrow (B_1 \cup \dots \cup B_m)$

The empty sequent

$$\longrightarrow$$

means a **contradiction**

Gentzen Sequents

Given **non empty** sequences Γ, Δ

We denote by σ_{Γ} any **conjunction** of all formulas of Γ

We denote by δ_{Δ} any **disjunction** of all formulas of Δ

The **intuitive semantics** of a non- empty sequent $\Gamma \longrightarrow \Delta$ is

$$\Gamma \longrightarrow \Delta \equiv (\sigma_{\Gamma} \Rightarrow \delta_{\Delta})$$

Formal Semantics

Formal semantics for **sequents** of **GL** is defined as follows

Let $v : VAR \rightarrow \{T, F\}$ be a truth assignment and v^* its extension to the set of formulas \mathcal{F}

We **extend** v^* to the set

$$SQ = \{ \Gamma \rightarrow \Delta : \Gamma, \Delta \in \mathcal{F}^* \}$$

of all sequents as follows

For any sequent $\Gamma \rightarrow \Delta \in SQ$

$$v^*(\Gamma \rightarrow \Delta) = v^*(\sigma_\Gamma) \Rightarrow v^*(\delta_\Delta)$$

Formal Semantics

In the case when $\Gamma = \emptyset$ or $\Delta = \emptyset$ we **define**

$$v^*(\longrightarrow \Delta) = (T \Rightarrow v^*(\delta_\Delta))$$

$$v^*(\Gamma \longrightarrow) = (v^*(\sigma_\Gamma) \Rightarrow F)$$

The sequent $\Gamma \longrightarrow \Delta$ is **satisfiable** if there is a truth assignment $v : \text{VAR} \longrightarrow \{T, F\}$ such that

$$v^*(\Gamma \longrightarrow \Delta) = T$$

Formal Semantics

Model for $\Gamma \rightarrow \Delta$ is any v such that

$$v^*(\Gamma \rightarrow \Delta) = T$$

We write it $v \models \Gamma \rightarrow \Delta$

Counter-model is any v such that

$$v^*(\Gamma \rightarrow \Delta) = F$$

We write it $v \not\models \Gamma \rightarrow \Delta$

Tautology is any sequent $\Gamma \rightarrow \Delta$ such that

$v^*(\Gamma \rightarrow \Delta) = T$ for all truth assignments $v : VAR \rightarrow \{T, F\}$

We write it

$$\models \Gamma \rightarrow \Delta$$

Example

Example

Let $\Gamma \rightarrow \Delta$ be a sequent

$$a, (b \cap a) \rightarrow \neg b, (b \Rightarrow a)$$

The truth assignment v for which

$$v(a) = T \quad \text{and} \quad v(b) = T$$

is a **model** for $\Gamma \rightarrow \Delta$ as shows the following computation

$$\begin{aligned} v^*(a, (b \cap a) \rightarrow \neg b, (b \Rightarrow a)) &= v^*(\sigma_{\{a, (b \cap a)\}}) \Rightarrow v^*(\delta_{\{\neg b, (b \Rightarrow a)\}}) \\ &= v(a) \cap (v(b) \cap v(a)) \Rightarrow \neg v(b) \cup (v(b) \Rightarrow v(a)) \\ &= T \cap T \cap T \Rightarrow \neg T \cup (T \Rightarrow T) = T \Rightarrow (F \cup T) = T \Rightarrow T = T \end{aligned}$$

Example

Observe that the truth assignment v for which

$$v(a) = T \quad \text{and} \quad v(b) = T$$

is the **only one** for which

$$v^*(\Gamma) = v^*(a, (b \cap a) = T$$

and we proved that it is a **model** for

$$a, (b \cap a) \longrightarrow \neg b, (b \Rightarrow a)$$

It is hence **impossible** to find v which would **falsify it**, what proves that

$$\models a, (b \cap a) \longrightarrow \neg b, (b \Rightarrow a)$$

Gentzen System **GL**

Definition of **GL**

Logical Axioms **LA**

We adopt as an **axiom** any sequent of **variables (positive literals)** which contains a propositional variable that appears on **both sides** of the sequent arrow \longrightarrow , i.e any sequent of the form

$$\Gamma'_1, a, \Gamma'_2 \longrightarrow \Delta'_1, a, \Delta'_2$$

for any $a \in \mathit{VAR}$ and any sequences $\Gamma'_1, \Gamma'_2, \Delta'_1, \Delta'_2 \in \mathit{VAR}^*$

Gentzen System **GL**

Inference rules of **GL**

Let $\Gamma', \Delta' \in VAR^*$ and $\Gamma, \Delta \in \mathcal{F}^*$

Conjunction rules

$$(\cap \rightarrow) \frac{\Gamma', A, B, \Gamma \rightarrow \Delta'}{\Gamma', (A \cap B), \Gamma \rightarrow \Delta'}$$

$$(\rightarrow \cap) \frac{\Gamma \rightarrow \Delta, A, \Delta' ; \Gamma \rightarrow \Delta, B, \Delta'}{\Gamma \rightarrow \Delta, (A \cap B), \Delta'}$$

Gentzen System **GL**

Disjunction rules

$$(\rightarrow \cup) \frac{\Gamma \rightarrow \Delta, A, B, \Delta'}{\Gamma \rightarrow \Delta, (A \cup B), \Delta'}$$

$$(\cup \rightarrow) \frac{\Gamma', A, \Gamma \rightarrow \Delta' ; \Gamma', B, \Gamma \rightarrow \Delta'}{\Gamma', (A \cup B), \Gamma \rightarrow \Delta'}$$

Gentzen System **GL**

Implication rules

$$(\rightarrow\Rightarrow) \frac{\Gamma', A, \Gamma \rightarrow \Delta, B, \Delta'}{\Gamma', \Gamma \rightarrow \Delta, (A \Rightarrow B), \Delta'}$$

$$(\Rightarrow\rightarrow) \frac{\Gamma', \Gamma \rightarrow \Delta, A, \Delta' ; \Gamma', B, \Gamma \rightarrow \Delta, \Delta'}{\Gamma', (A \Rightarrow B), \Gamma \rightarrow \Delta, \Delta'}$$

Gentzen System **GL**

Negation rules

$$(\neg \rightarrow) \frac{\Gamma', \Gamma \rightarrow \Delta, A, \Delta'}{\Gamma', \neg A, \Gamma \rightarrow \Delta, \Delta'}$$

$$(\rightarrow \neg) \frac{\Gamma', A, \Gamma \rightarrow \Delta, \Delta'}{\Gamma', \Gamma \rightarrow \Delta, \neg A, \Delta'}$$

Gentzen System **GL**

We define the Gentzen System **GL**

$$\mathbf{GL} = (\mathcal{L}_{\{\cup, \cap, \Rightarrow, \neg\}}, \mathbf{SQ}, \mathbf{LA}, \mathcal{R})$$

for

$$\mathcal{R} = \{ (\cap \rightarrow), (\rightarrow \cap), (\cup \rightarrow), (\rightarrow \cup), (\Rightarrow \rightarrow), (\rightarrow \Rightarrow) \} \\ \cup \{ (\neg \rightarrow), (\rightarrow \neg) \}$$

We write, as usual,

$$\vdash_{\mathbf{GL}} \Gamma \rightarrow \Delta$$

to denote that $\Gamma \rightarrow \Delta$ has a formal proof in **GL**

A formula $A \in \mathcal{F}$, has a proof in **GL** if the sequent $\rightarrow A$ has a proof in **GL**, i.e.

$$\vdash_{\mathbf{GL}} A \quad \text{if and only if} \quad \rightarrow A$$

Gentzen System **GL**

We consider, as we did with **RS** the proof trees for **GL**, i.e. we define

A **proof tree**, or **GL**-proof of $\Gamma \longrightarrow \Delta$ is a tree

$$\mathbf{T}_{\Gamma \longrightarrow \Delta}$$

of sequents satisfying the following conditions:

1. The topmost sequent, i.e **the root** of $\mathbf{T}_{\Gamma \longrightarrow \Delta}$ is $\Gamma \longrightarrow \Delta$
2. All **leafs** are **axioms**
3. The **nodes** are sequents such that each sequent on the tree follows from the ones immediately preceding it by one of the rules.

Gentzen System **GL**

Remark

The **proof search** in **GL** as defined by the **decomposition tree** for a given formula **A is not always unique**

We show it on an example on the next slide

Example

A tree-proof in **GL** of the de Morgan Law

$$\longrightarrow (\neg(a \wedge b) \Rightarrow (\neg a \vee \neg b))$$

$$| (\longrightarrow \Rightarrow)$$

$$\neg(a \wedge b) \longrightarrow (\neg a \vee \neg b)$$

$$| (\longrightarrow \vee)$$

$$\neg(a \wedge b) \longrightarrow \neg a, \neg b$$

$$| (\longrightarrow \neg)$$

$$b, \neg(a \wedge b) \longrightarrow \neg a$$

$$| (\longrightarrow \neg)$$

$$b, a, \neg(a \wedge b) \longrightarrow$$

$$| (\neg \longrightarrow)$$

$$b, a \longrightarrow (a \wedge b)$$

$$\bigwedge (\longrightarrow \cap)$$

$$b, a \longrightarrow a$$

$$b, a \longrightarrow b$$

Example

Here is another tree-proof in **GL** of the de Morgan Law

$$\rightarrow (\neg(a \cap b) \Rightarrow (\neg a \cup \neg b))$$

$$| (\rightarrow \Rightarrow)$$

$$\neg(a \cap b) \rightarrow (\neg a \cup \neg b)$$

$$| (\rightarrow \cup)$$

$$\neg(a \cap b) \rightarrow \neg a, \neg b$$

$$| (\rightarrow \neg)$$

$$b, \neg(a \cap b) \rightarrow \neg a$$

$$| (\neg \rightarrow)$$

$$b \rightarrow \neg a, (a \cap b)$$

$$\bigwedge (\rightarrow \cap)$$

$$b \rightarrow \neg a, a$$

$$| (\rightarrow \neg)$$

$$b, a \rightarrow a$$

$$b \rightarrow \neg a, b$$

$$| (\rightarrow \neg)$$

$$b, a \rightarrow b$$

Gentzen System **GL**

The process of **searching for proofs** of a formula **A** in **GL** consists, as in the **RS** type systems, of building certain trees, called **decomposition trees**

Their construction is similar to the one for **RS** type systems

We take a **root** of a **decomposition tree** T_A of of a formula **A**
a sequent $\longrightarrow A$

For each **node**, if there is a **rule** of **GL** which conclusion has the same form as **node** sequent, then the **node** has **children** that are **premises** of the **rule**

If the **node** consists only of a sequent built only out of **variables** then it **does not** have any **children**

This is a **termination condition** for the **tree**

Gentzen System **GL**

We **prove** that each formula **A** generates a **finite** set of decomposition trees, \mathcal{T}_A , such that the following holds

If there exist a tree $T_A \in \mathcal{T}_A$ whose **all leafs** are **axioms**, then tree T_A constitutes a **proof** of **A** in **GL**

If **all trees** in \mathcal{T}_A have at **least one non-axiom leaf**, the proof of **A** **does not exist**

The **first step** in **defining** a notion of a **decomposition tree** consists of **transforming** the inference rules of **GL**, as we did in the case of the **RS** type systems, into corresponding **decomposition rules**

Decomposition Rules of **GL**

Decomposition rules of **GL**

Let $\Gamma', \Delta' \in VAR^*$ and $\Gamma, \Delta \in \mathcal{F}^*$

Conjunction rules

$$(\cap \rightarrow) \frac{\Gamma', (A \cap B), \Gamma \rightarrow \Delta'}{\Gamma', A, B, \Gamma \rightarrow \Delta'}$$

$$(\rightarrow \cap) \frac{\Gamma \rightarrow \Delta, (A \cap B) \Delta'}{\Gamma \rightarrow \Delta, A, \Delta' ; \Gamma \rightarrow \Delta, B, \Delta'}$$

Decomposition Rules of **GL**

Disjunction rules

$$(\rightarrow \cup) \frac{\Gamma \rightarrow \Delta, (A \cup B), \Delta'}{\Gamma \rightarrow \Delta, A, B, \Delta'}$$

$$(\cup \rightarrow) \frac{\Gamma', (A \cup B), \Gamma \rightarrow \Delta'}{\Gamma', A, \Gamma \rightarrow \Delta' ; \Gamma', B, \Gamma \rightarrow \Delta'}$$

Decomposition Rules of **GL**

Implication rules

$$(\rightarrow\Rightarrow) \frac{\Gamma', \Gamma \rightarrow \Delta, (A \Rightarrow B), \Delta'}{\Gamma', A, \Gamma \rightarrow \Delta, B, \Delta'}$$

$$(\Rightarrow\rightarrow) \frac{\Gamma', (A \Rightarrow B), \Gamma \rightarrow \Delta, \Delta'}{\Gamma', \Gamma \rightarrow \Delta, A, \Delta' ; \Gamma', B, \Gamma \rightarrow \Delta, \Delta'}$$

Decomposition Rules of **GL**

Negation rules

$$(\neg \rightarrow) \frac{\Gamma', \neg A, \Gamma \rightarrow \Delta, \Delta'}{\Gamma', \Gamma \rightarrow \Delta, A, \Delta'}$$

$$(\rightarrow \neg) \frac{\Gamma', \Gamma \rightarrow \Delta, \neg A, \Delta'}{\Gamma', A, \Gamma \rightarrow \Delta, \Delta'}$$

Decomposition Tree Definition

For each formula $A \in \mathcal{F}$, a decomposition tree T_A is a tree build as follows

Step 1. The sequent $\longrightarrow A$ is the **root** of T_A

For any node $\Gamma \longrightarrow \Delta$ of the tree we follow the steps below

Step 2. If $\Gamma \longrightarrow \Delta$ is **indecomposable**, then $\Gamma \longrightarrow \Delta$ becomes a **leaf** of the tree

Step 3. If $\Gamma \longrightarrow \Delta$ is **decomposable**, then we pick a **decomposition rule** that **maches** the sequent of the **current node**

To do so we **proceed** as follows

Decomposition Tree Definition

1 Given a node $\Gamma \rightarrow \Delta$

We traverse Γ from **left** to **right** to find the **first decomposable formula**

Its **main connective**. \circ identifies a **possible decomposition rule** ($\circ \rightarrow$) Then we **check** if this decomposition rule ($\circ \rightarrow$) applies. If it does we put its **conclusions** (conclusion) as **leaves** (leaf)

2 We **traverse** Δ from **right** to **left** to find the **first decomposable formula**

Its **main connective** \circ identifies a **possible decomposition rule** ($\rightarrow \circ$)

Then we check if this decomposition rule applies. If it does we put its **conclusions** (conclusion) as **leaves** (leaf)

3 If **1** and **2** **applies** we **choose one of the rules**

Step 4. We repeat **Step 2.** and **Step 3.** until we obtain **only leaves**

Decomposition Tree Definition

Observe that a decomposable $\Gamma \longrightarrow \Delta$ is always in the domain in one of the decomposition rules $(\circ \longrightarrow)$, $(\longrightarrow \circ)$, or in the domain of both. Hence the tree \mathbf{T}_A may not be unique and all possible choices of 3. give all possible decomposition trees

System **GL** Exercises

Exercise

Prove, by constructing a proper **decomposition tree** that

$$\vdash_{\mathbf{GL}} ((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a))$$

Solution

By definition, we have that

$$\vdash_{\mathbf{GL}} ((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a)) \text{ if and only if}$$

$$\vdash_{\mathbf{GL}} \longrightarrow ((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a))$$

We construct a decomposition tree $\mathbf{T}_{\rightarrow A}$ as follows

System **GL** Exercises

T \rightarrow **A**

$\rightarrow ((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a))$

| $(\rightarrow \Rightarrow)$

$(\neg a \Rightarrow b) \rightarrow (\neg b \Rightarrow a)$

| $(\rightarrow \Rightarrow)$

$\neg b, (\neg a \Rightarrow b) \rightarrow a$

| $(\rightarrow \neg)$

$(\neg a \Rightarrow b) \rightarrow b, a$

$\bigwedge (\Rightarrow \rightarrow)$

$\rightarrow \neg b, b, a$

| $(\rightarrow \neg)$

$b \rightarrow b, a$

axiom

$b \rightarrow b, a$

axiom

All leaves of the tree are **axioms**, hence we have found the **proof** of **A** in **GL**

System **GL** Exercises

Exercise

Prove, by constructing proper **decomposition trees** that

$$\not\vdash_{\mathbf{GL}} ((a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a))$$

Solution

Observe that for any formula A , its decomposition tree $T_{\rightarrow A}$ in **GL** may **not be unique**

Hence we have to construct **all possible decomposition trees** to prove that **each** of them has a **non axiom leaf**

We construct the decomposition trees for $\rightarrow A$ as follows

System **GL** Exercises

T_{1→A}

→ ((a ⇒ b) ⇒ (¬b ⇒ a))

| (→⇒) (*one choice*)

(a ⇒ b) → (¬b ⇒ a)

| (→⇒) (*first of two choices*)

¬b, (a ⇒ b) → a

| (¬→) (*one choice*)

(a ⇒ b) → b, a

∧ (⇒→) (*one choice*)

→ a, b, a

non - axiom

b → b, a

axiom

The tree contains a **non- axiom** leaf, hence it is **not a proof**

We have **one more tree** to construct

System **GL** Exercises

T_{1→A}

$$\rightarrow ((a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a))$$

$$| (\rightarrow \Rightarrow) \text{ (one choice)}$$

$$(a \Rightarrow b) \rightarrow (\neg b \Rightarrow a)$$

$$\wedge (\Rightarrow \rightarrow) \text{ (second choice)}$$

$$\rightarrow (\neg b \Rightarrow a), a$$

$$| (\rightarrow \Rightarrow) \text{ (one choice)}$$

$$\neg b \rightarrow a, a$$

$$| (\neg \rightarrow) \text{ (one choice)}$$

$$\rightarrow a, a, b$$

non - axiom

$$b \rightarrow (\neg b \Rightarrow a)$$

$$| (\rightarrow \Rightarrow) \text{ (one choice)}$$

$$b, \neg b \rightarrow a$$

$$| (\neg \rightarrow) \text{ (one choice)}$$

$$b \rightarrow a, b$$

axiom

All possible trees end with a **non-axiom leaf**. It proves that

$$\not\vdash_{\text{GL}} ((a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a))$$

System **GL** Exercises

Does the tree below constitute a proof in **GL** ? Justify your answer

$$\begin{array}{c} \mathbf{T}_{\rightarrow A} \\ \rightarrow \neg\neg((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a)) \\ \quad | (\rightarrow \neg) \\ \neg((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a)) \rightarrow \\ \quad | (\neg \rightarrow) \\ \rightarrow ((\neg a \Rightarrow b) \Rightarrow (\neg b \Rightarrow a)) \\ \quad | (\rightarrow \Rightarrow) \\ (\neg a \Rightarrow b) \rightarrow (\neg b \Rightarrow a) \\ \quad | (\rightarrow \Rightarrow) \\ (\neg a \Rightarrow b), \neg b \rightarrow a \\ \quad | (\neg \rightarrow) \\ (\neg a \Rightarrow b) \rightarrow b, a \\ \quad \bigwedge (\Rightarrow \rightarrow) \end{array}$$

$$\begin{array}{cc} \rightarrow \neg a, b, a & b \rightarrow b, a \\ | (\rightarrow \neg) & \text{axiom} \\ a \rightarrow b, a & \\ \text{axiom} & \end{array}$$

System **GL** Exercises

Solution

The tree $\mathbf{T}_{\rightarrow A}$ is **not a proof** in **GL** because a rule corresponding to the **decomposition step** below **does not exist** in **GL**

$$(\neg a \Rightarrow b), \neg b \longrightarrow a$$

$$| (\neg \rightarrow)$$

$$(\neg a \Rightarrow b) \longrightarrow b, a$$

It is a proof in some system **GL1** that has all the rules of **GL** except its rule $(\neg \rightarrow)$

$$(\neg \rightarrow) \quad \frac{\Gamma', \Gamma \longrightarrow \Delta, A, \Delta'}{\Gamma', \neg A, \Gamma \longrightarrow \Delta, \Delta'}$$

This rule has to be replaced in by the rule:

$$(\neg \rightarrow)_1 \quad \frac{\Gamma, \Gamma' \longrightarrow \Delta, A, \Delta'}{\Gamma, \neg A, \Gamma' \longrightarrow \Delta, \Delta'}$$

Exercises

Exercise 1

Write **all proofs** in **GL** of

$$(\neg(a \wedge b) \Rightarrow (\neg a \vee \neg b))$$

Exercise 2

Find a formula which has a **unique** decomposition tree

Exercise 3

Describe for which kind of formulas the decomposition tree is unique

System **GL** Exercises

Exercise

We know that the system **GL** is **strongly sound**

Prove, by constructing a **counter-model determined** by a proper **decomposition tree** that

$$\not\models ((b \Rightarrow a) \Rightarrow (\neg b \Rightarrow a))$$

We construct the decomposition tree for the formula

$A : ((b \Rightarrow a) \Rightarrow (\neg b \Rightarrow a))$ as follows

System **GL** Exercises

T_{→A}

→ ((b ⇒ a) ⇒ (¬b ⇒ a))

| (⇒)

(b ⇒ a) → (¬b ⇒ a)

| (⇒)

¬b, (b ⇒ a) → a

| (¬ →)

(b ⇒ a) → b, a

∧ (⇒ →)

→ b, b, a

non - axiom

a → b, a

axiom

The **counter model** determined by the tree **T**_{→A} is any truth assignment **v** that evaluates the **non axiom leaf** → b, b, a to **F**

System **GL** Exercises

Let $v : VAR \rightarrow \{T, F\}$ be a truth assignment

By definition of semantic for sequents we have that

$$v^*(\rightarrow b, b, a) = (T \Rightarrow v(b) \cup v(b) \cup v(a))$$

Hence $v^*(\rightarrow b, b, a) = F$ if and only if

$$(T \Rightarrow v(b) \cup v(b) \cup v(a)) = F \text{ if and only if}$$
$$v(b) = v(a) = F$$

The **counter model** determined by the $\mathbf{T}_{\rightarrow A}$ is any

$$v : VAR \rightarrow \{T, F\} \text{ such that } v(b) = v(a) = F$$

System **GL** Exercises

Exercise

Prove the **Completeness theorem** for **GL**

Assume that the **Strong Soundness** has been already proved and the **Decompositions Trees** are already defined

Reminder

Formula Completeness for **GL**: For any $A \in \mathcal{F}$,

$$\models A \text{ if and only if } \vdash_{GL} \rightarrow A$$

Soundness for **GL**: For any $A \in \mathcal{F}$,

$$\text{If } \vdash_{GL} \rightarrow A, \text{ then } \models A$$

Completeness part for **GL**: For any $A \in \mathcal{F}$,

$$\text{If } \models A, \text{ then } \vdash_{GL} \rightarrow A$$

Proof of Completeness of **GL**

We prove the logically equivalent form of the **Completeness part**: For any $A \in \mathcal{F}$,

If $\not\vdash_{GL} \rightarrow A$ then $\not\models A$

Proof

Assume $\not\vdash_{GL} \rightarrow A$, i.e. $\rightarrow A$ does not have a proof in **GL**

Let \mathcal{T}_A be a **set of all decomposition trees** of $\rightarrow A$

As $\not\vdash_{GL} \rightarrow A$, each $T \in \mathcal{T}_A$ has a **non-axiom leaf**

We choose an arbitrary $T_A \in \mathcal{T}_A$

Proof of Completeness of GL

Let $\Gamma' \rightarrow \Delta', \Gamma', \Delta' \in VAR^*$ be the **non-axiom leaf** of the tree T_A

The non-axiom leaf $\Gamma' \rightarrow \Delta'$ **determines** a truth assignment $v : VAR \rightarrow \{T, F\}$ which is defined as follows:

$$v(a) = \begin{cases} T & \text{if } a \text{ appears in } \Gamma' \\ F & \text{if } a \text{ appears in } \Delta' \\ \text{any value} & \text{if } a \text{ does not appear in } \Gamma' \rightarrow \Delta' \end{cases}$$

By the **strong soundness** of the rules of inference of **GL** it proves that $v^*(A) = F$, i.e. that $\not\models A$

Original Gentzen systems
LK for Classical Propositional Logic and **LI** for Intuitionistic Logic

Gentzen Systems **LK**, **LI** for Classical and Intuitionistic Logics

The proof systems **LK** for **Classical** Propositional Logic and **LI** for **Intuitionistic** Propositional Logic as presented here were published by **G. Gentzen** in **1935**

The proof system **LI** for **Intuitionistic Logic** was presented as a **particular case** of his proof system **LK** for the **classical logic**

Classical Gentzen System **LK**

Language of **LK**

$$\mathcal{L} = \mathcal{L}_{\{\neg, \wedge, \vee, \Rightarrow\}} \quad \text{and} \quad \mathcal{E} = \text{SQ}$$

for

$$\text{SQ} = \{\Gamma \longrightarrow \Delta : \Gamma, \Delta \in \mathcal{F}^*\}$$

Axioms of LK any sequent of the form

$$\Gamma_1, A, \Gamma_2 \longrightarrow \Gamma_3, A, \Gamma_4$$

Classical Gentzen System **LK**

Rules of inference of **LK** are as follows

Structural Rules

Weakening

$$(weak \rightarrow) \frac{\Gamma \rightarrow \Delta}{A, \Gamma \rightarrow \Delta}$$

$$(\rightarrow weak) \frac{\Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta, A}$$

Contraction

$$(contr \rightarrow) \frac{A, A, \Gamma \rightarrow \Delta}{A, \Gamma \rightarrow \Delta}$$

$$(\rightarrow contr) \frac{\Gamma \rightarrow \Delta, A, A}{\Gamma \rightarrow \Delta, A}$$

Classical Gentzen System **LK**

Structural Rules

Exchange

$$(exch \rightarrow) \frac{\Gamma_1, A, B, \Gamma_2 \rightarrow \Delta}{\Gamma_1, B, A, \Gamma_2 \rightarrow \Delta}$$

$$(\rightarrow exch) \frac{\Delta \rightarrow \Gamma_1, A, B, \Gamma_2}{\Delta \rightarrow \Gamma_1, B, A, \Gamma_2}$$

Classical Gentzen System **LK**

Logical Rules

Conjunction rules

$$(\cap \rightarrow) \frac{A, B, \Gamma \rightarrow \Delta}{(A \cap B), \Gamma \rightarrow \Delta}$$

$$(\rightarrow \cap) \frac{\Gamma \rightarrow \Delta, A \quad ; \quad \Gamma \rightarrow \Delta, B, \Delta}{\Gamma \rightarrow \Delta, (A \cap B)}$$

Disjunction rules

$$(\rightarrow \cup) \frac{\Gamma \rightarrow \Delta, A, B}{\Gamma \rightarrow \Delta, (A \cup B)}$$

$$(\cup \rightarrow) \frac{A, \Gamma \rightarrow \Delta \quad ; \quad B, \Gamma \rightarrow \Delta}{(A \cup B), \Gamma \rightarrow \Delta}$$

Classical Gentzen System **LK**

Implication rules

$$(\rightarrow\Rightarrow) \quad \frac{A, \Gamma \rightarrow \Delta, B}{\Gamma \rightarrow \Delta, (A \Rightarrow B)}$$

$$(\Rightarrow\rightarrow) \quad \frac{\Gamma \rightarrow \Delta, A \quad ; \quad B, \Gamma \rightarrow \Delta}{(A \Rightarrow B), \Gamma \rightarrow \Delta}$$

Classical Gentzen System **LK**

Negation rules

$$(\neg \rightarrow) \frac{\Gamma \rightarrow \Delta, A}{\neg A, \Gamma \rightarrow \Delta}$$

$$(\rightarrow \neg) \frac{A, \Gamma \rightarrow \Delta}{\Gamma \rightarrow \Delta, \neg A}$$

We define formally

LK = (\mathcal{L} , SQ, AX, Structural rules, Logical rules)

Intuitionistic I Gentzen System **LI**

Language of **LI**

Any expression

$$\Gamma \longrightarrow \Delta$$

where $\Gamma, \Delta \in \mathcal{F}^*$ and

Δ consists of **at most one formula**

is called a **LI sequent**

We denote the set of all **LI sequents** by *ISQ*, i.e.

$$ISQ = \{\Gamma \longrightarrow \Delta : \Delta \text{ consists of } \mathbf{at\ most\ one\ formula}\}$$

Axioms of LI

Axioms of LI consist of any sequent from the set *ISQ* which contains a **formula** that appears on **both sides** of the sequent arrow \longrightarrow , i.e any sequent of the form

$$\Gamma, A, \Delta \longrightarrow A$$

for $\Gamma, \Delta \in \mathcal{F}^*$

Rules of Inference of LI

The set inference rules of LI is divided into **two groups** : the **structural rules** and the **logical rules**

There are three **Structural Rules** of LI: **Weakening**, **Contraction** and **Exchange**

Weakening structural rule

$$(weak \rightarrow) \frac{\Gamma \rightarrow \Delta}{A, \Gamma \rightarrow \Delta}$$

$$(\rightarrow weak) \frac{\Gamma \rightarrow}{\Gamma \rightarrow A}$$

A is called the **weakening formula**

Remember that Δ contains **at most one formula**

Rules of Inference of **LI**

Contraction structural rule

$$(contr \rightarrow) \frac{A, A, \Gamma \rightarrow \Delta}{A, \Gamma \rightarrow \Delta}$$

The case below is **not VALID** for **LI**; we list it as it will be used in the classical case

$$(\rightarrow contr) \frac{\Gamma \rightarrow \Delta, A, A}{\Gamma \rightarrow \Delta, A}$$

A is called the **contraction formula**

Remember that Δ contains **at most one formula**

Rules of Inference of **LI**

Exchange structural rule

$$(exch \rightarrow) \frac{\Gamma_1, A, B, \Gamma_2 \rightarrow \Delta}{\Gamma_1, B, A, \Gamma_2 \rightarrow \Delta}$$

The case below is **not VALID** for **LI**; we list it as it will be used in the classical case

$$(\rightarrow exch) \frac{\Delta \rightarrow \Gamma_1, A, B, \Gamma_2}{\Delta \rightarrow \Gamma_1, B, A, \Gamma_2}.$$

Remember that Δ contains **at most one formula**

Rules of Inference of LI

Logical Rules

Conjunction rules

$$(\wedge \rightarrow) \frac{A, B, \Gamma \rightarrow \Delta}{(A \wedge B), \Gamma \rightarrow \Delta},$$

$$(\rightarrow \wedge) \frac{\Gamma \rightarrow A ; \Gamma \rightarrow B}{\Gamma \rightarrow (A \wedge B)}$$

Remember that Δ contains **at most one formula**

Rules of Inference of LI

Disjunction rules

$$(\rightarrow \cup)_1 \quad \frac{\Gamma \rightarrow A}{\Gamma \rightarrow (A \cup B)}$$

$$(\rightarrow \cup)_2 \quad \frac{\Gamma \rightarrow B}{\Gamma \rightarrow (A \cup B)}$$

$$(\cup \rightarrow) \quad \frac{A, \Gamma \rightarrow \Delta ; B, \Gamma \rightarrow \Delta}{(A \cup B), \Gamma \rightarrow \Delta}$$

Remember that Δ contains **at most one formula**