

**CSE541 INTRODUCTION EXERCISES on SETS  
SOLUTIONS**

**QUESTION 1** Use the above definition to prove the following

**FACT 1** A set  $A$  is INFINITE iff it contains a countably infinite subset, i.e. one can define a 1-1 sequence  $\{a_n\}_{n \in \mathbb{N}}$  of some elements of  $A$ .

**SOLUTION 1.** Implication  $\rightarrow$

If  $A$  is infinite, then we can define a 1-1 sequence of elements of  $A$ .

Let  $A$  be infinite,

We define a sequence

$$a_1, \dots, a_n, \dots$$

as follows.

1. Observe that  $A \neq \emptyset$ , because if  $A = \emptyset$ ,  $A$  would be finite. contradiction.

So there is an element of  $a \in A$ .

We define

$$a_1 = a$$

2. Consider a set  $A - \{a_1\} = A_1$ .  $A_1 \neq \emptyset$  because if  $A_1 = \emptyset$ , then  $A - \{a_1\} = \emptyset$  and  $A$  is finite. Contradiction.

So there is an element  $a_2 \in A - \{a_1\}$  and  $a_1 \neq a_2$ .

We defined

$$a_1, a_2$$

3. Assume now that we have defined an  $n$ -elements and sequence

$$a_1, a_2, \dots, a_n \text{ for } a_1 \neq a_2 \neq \dots \neq a_n$$

Consider a set  $A_n = A - \{a_1, \dots, a_n\}$ .

The set  $A_n \neq \emptyset$  because if  $A - \{a_1, \dots, a_n\} = \emptyset$ , then  $A$  is finite. Contradiction

So there is an element

$$a_{n+1} \in A - \{a_1, \dots, a_n\}$$

and  $a_{n+1} \neq a_n \neq \dots \neq a_1$

By mathematical induction,

we have defined a 1-1 sequence

$$a_1, a_2, \dots, a_n, \dots$$

elements of  $A$ .

2. Implication  $\leftarrow$

If  $A$  contain a 1-1 sequence, then  $A$  is infinite.

Assume  $A$  is not infinite; i.e  $A$  is finite. Every subset of finite set is finite, so we can't have a 1-1 infinite sequence of elements of  $A$ . Contradiction.




Figure 1: problem 1

**QUESTION 2** Use the above definition and FACT 1 from Question 1 to prove the following characterization of infinite sets.

**Dedekind Theorem** A set  $A$  is INFINITE iff there is a set proper subset  $B$  of the set  $A$  such that  $|A| = |B|$ .

**SOLUTION** Part1. If  $A$  is infinite, then there is  $B \subsetneq A$  and

$$f : A \xrightarrow[\text{onto}]{1-1} B$$

$A$  is infinite, by Q1, we have a 1-1 sequence

$$a_1, a_2, \dots, a_n, \dots$$

of elements  $A$ .

We take  $B = A - \{a_1\}$ ,  $B \subsetneq A$  and we define a function

$$f : A \xrightarrow[\text{onto}]{1-1} B$$

as follows

$$f(a_1) = a_2$$

$$f(a_2) = a_3$$

⋮

$$f(a_n) = a_{n+1}$$

$$f(a) = a, \text{ for all other } a \in A$$




Figure 2: problem 2:part 1

obviously,  $f$  is 1-1, onto

Observe: we have other choices of B!

Part 2. Assume that we have  $B \subsetneq A$  are

$$f : A \xrightarrow[\text{onto}]{1-1} B$$

We use Q1 to show that A is infinite; i.e we construct an 1-1 sequence  $a_1 \dots a_n$  of elements of  $A_n$  as follows.

$B \subsetneq A$ , so  $A - B \neq \emptyset$  and we have  $b \in A - B$ . This is our first element of the sequence.

Observe:  $f : A \xrightarrow[\text{onto}]{1-1} B$ , so  $f(b) \in B$  and  $b \in A - B$ , hence  $f(b) \neq b$  and  $f(b)$  is our second element of the sequence.

We have now,

$$b, f(b) \quad f(b) \neq b, b \in A - B, f(b) \in B$$

Take new,

$ff(b)$ . As  $f$  is 1-1 and  $f(b) \neq b$ , we get  $ff(b) \neq f(b) \neq b$ ,  $ff(b) \in B$  and the sequence  $b, f(b), ff(b)$  is 1-1.

We create  $ff(b) = f^2(b)$

We continue the construction by mathematical induction.

Assume that we have constructed a 1-1 sequence

$$b, f(b), f^2(b), f^3(b), \dots, f^n(b)$$

Observe that  $ff^n(b) = f^{n+1}(b) \neq f^n(b)$  as  $f$  is 1-1.

By mathematical induction, we have that  $\{f^n(b)\}_{n \in \mathbb{N}}$  is a 1-1 sequence of elements of A and hence A is infinite.




Figure 3: problem 2:part 2

**QUESTION 3** Use technique from DEDEKIND THEOREM to prove the following

**Theorem** For any infinite set  $A$  and its finite subset  $B$ ,  $|A| = |A - B|$ .

**SOLUTION**  $A$  is infinite, then by Q1 there is a 1-1 sequence:

$$a_1, a_2, \dots, a_n, \dots$$

of elements of  $A$ .

Let  $|B| = K$ . We choose  $K$  1-1 sequence  $\{C_n^k\}_{n \in \mathbb{N}}$  of the sequence  $\{a_n\}_{n \in \mathbb{N}}$ .

Let  $B = \{b_1, \dots, b_k\}$ . We construct a function  $f : A \xrightarrow[\text{onto}]{1-1} A - \{b_1, \dots, b_k\}$  as follows

$$\begin{aligned} f(b_1) &= c_1^1, & f(c_1^1) &= c_2^1, \dots, f(c_n^1) &= c_{n+1}^1 \\ f(b_2) &= c_1^2, & f(c_1^2) &= c_2^2, \dots, f(c_n^2) &= c_{n+1}^2 \\ & & & \vdots & \\ f(b_k) &= c_1^k, & f(c_1^k) &= c_2^k, \dots, f(c_n^k) &= c_{n+1}^k \\ f(a) &= a \text{ all } a \in A - B \end{aligned}$$

As all sequences  $\{C_n^m\}_{n \in \mathbb{N}, m=1, \dots, k}$  are 1-1, the function  $f$  is 1-1 and obviously ONTO  $A - B$ .

**QUESTION 4** Use DEDEKIND THEOREM to prove that the set  $\mathbb{N}$  of natural numbers is infinite.

**SOLUTION** We use Dedekind theorem i.e we must define  $f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} B \subsetneq \mathbb{N}$ . There are many such function

for example  $f(n) = n + 1, f : \mathbb{N} \xrightarrow[\text{onto}]{1-1} \mathbb{N} - \{0\}$

One can also use Q1 and define any 1-1 sequences in  $\mathbb{N}$ .

**QUESTION 5** Use DEDEKIND THEOREM to prove that the set  $\mathbb{R}$  of real numbers is infinite.




Figure 4: problem 3

**SOLUTION** We use Dedekind theorem

$$f(x) = 2^x \quad x \in \mathbb{R}$$

$$f : \mathbb{R} \xrightarrow[\text{onto}]{1-1} \mathbb{R}^+$$

One can also use Q1 and define any 1-1 sequences in  $\mathbb{R}$ .

**QUESTION 6** Use technique from DEDEKIND THEOREM to prove that the interval  $[a, b], a < b$  of real numbers is infinite and that  $|[a, b]| = |(a, b)|$ .

**SOLUTION1** Use construction in the proof of Q3.

$$f : [a, b] \xrightarrow[\text{onto}]{1-1} [a, b] - \{a, b\} = (a, b)$$

This is the solution I had in mine!

**SOLUTION2** Use Q3  $(a, b) = [a, b] - B, B$  :finite

**QUESTION 7** Prove, using the above definitions 3 and 4 that for any cardinal numbers  $\mathcal{M}, \mathcal{N}, \mathcal{K}$  the following formulas hold:

1.  $\mathcal{N} \leq \mathcal{N}$
2. If  $\mathcal{N} \leq \mathcal{M}$  and  $\mathcal{M} \leq \mathcal{K}$ , then  $\mathcal{N} \leq \mathcal{K}$ .

**SOLUTION** 1.  $\mathcal{N} \leq \mathcal{N}$  means that for any set  $A, |A| \leq |A|$

1.  $f(a) = a$  establishes  $f : A \xrightarrow{1-1} A$
2.  $\mathcal{N} \leq \mathcal{M}$  and  $\mathcal{M} \leq \mathcal{K}$ , then  $\mathcal{N} \leq \mathcal{K}$ .

We have  $|A| = \mathcal{N}, |B| = \mathcal{M}, |C| = \mathcal{K}$  and  $f : A \xrightarrow{1-1} B$  and  $g : B \xrightarrow{1-1} C$ , then we have to construct  $h : A \xrightarrow{1-1} C$ .

$h$  is a composition of  $f$  and  $g$ . i.e  $h(a) = g(f(a))$ , all  $a \in A$

**QUESTION 8** Prove, for any sets  $A, B, C$  the following holds.

**Fact 2**

*If  $C \subseteq B \subseteq A$  and  $|A| = |C|$ , then  $|A| = |B| = |C|$ .*

To prove  $|A| = |B|$  you must use definition 3, i.e to construct a proper function. Use the construction from proofs of Fact 1 and Question 3

**SOLUTION 1.**  $A, B, C$  are finite and  $|A| = |C|$ , and  $C \subseteq B \subseteq A$ , so  $A = B = C$ , and have  $|A| = |B| = |C|$

2.  $A, B, C$  are infinite sets, we have  $|A| = |C|$  i.e we have  $f : A \xrightarrow[\text{onto}]{1-1} C$

We want to construct a function

$$g : A \xrightarrow[\text{onto}]{1-1} B, \text{ where } A \subseteq B \subseteq C$$

Take  $A - B$ . We assume that  $A - B \neq \emptyset$ , if not,  $A = B$ , and  $|A| = |C|$  given  $|A| = |B| = |C|$ .

We consider case  $C \subset B \subset A$ . Take any  $a \in (A - B)$ , as  $f : A \xrightarrow[\text{onto}]{1-1} C$ ,  $f(a) \in C$ ,  $f$  is 1-1 so  $ff(a) \neq f(a)$

... in general  $f^n(a) \neq f^{n+1}(a)$  and we have a sequence for any  $a \in A - B$

$f(a), f^2(a), \dots, f^n(a) \dots$  of elements of  $C$ .

We construct a function  $g : A \xrightarrow[\text{onto}]{1-1} B$

$$g(a) = f(a)$$

$$g(f(a)) = f^2(a)$$

$$g(f^2(a)) = f^3(a)$$

⋮

$$g(f^n(a)) = f^{n+1}(a)$$

$$g(x) = x \quad \text{for all other } x \in A$$




Figure 5: problem 8: Figure of function  $g : A \xrightarrow[\text{onto}]{1-1} B$ .  $a, b$  represent any two element of  $A$

**QUESTION 9** Prove the following

**Berstein Theorem** (1898) For any cardinal numbers  $\mathcal{M}, \mathcal{N}$

$$\mathcal{N} \leq \mathcal{M} \text{ and } \mathcal{M} \leq \mathcal{N} \text{ then } \mathcal{N} = \mathcal{M}.$$

**SOLUTION** Let  $A, B$  be two sets such that  $|A| = \mathcal{N}, |B| = \mathcal{M}$ , we rewrite on theorem as

**Berstein Theorem** For any sets  $A, B$

If  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$

**case1.** The sets  $A, B$  are disjoint.

As  $|A| \leq |B|$ , we have a function  $f : A \xrightarrow{1-1} B$ , i.e  $f : A \xrightarrow[onto]{1-1} fA \subseteq B$  and  $|A| = |fA|$  where  $fA$  denotes the image of  $A$  under  $f$ .

As  $|B| \leq |A|$ , we have a function  $g : B \xrightarrow[onto]{1-1} gB \subseteq A$  and  $|B| = |gB|$

We picture it as follow.

$$|B| = |gB|, |A| = |fA|$$




Figure 6: problem 9

As  $f : A \xrightarrow{1-1} B$  and  $gB \subseteq A$ , we get  $fgB \subseteq fA$  and hence

$$fgB \subseteq fA \subseteq A \tag{1}$$

Also,  $gB \subseteq A$  and  $g : B \xrightarrow{1-1} B$ . Hence,  $fg : B \xrightarrow[onto]{1-1} fgB$  and

$$|B| = |fgB| \tag{2}$$

We have a following picture.




Figure 7: problem 9

By eq.2,  $|B| = |fgB|$  and by eq.1,  $fgB \subseteq fA \subseteq B$  and  $|B| = |fA|$

By Q8, we get

$$|fA| = |B|$$

Hence,  $|B| = |A|$

**case2.** the set  $A, B$  are NOT disjoint.

Repeat the same(or Google the proof) for the following picture.




Figure 8: problem 9