

## CSE547 CARDINALITIES OF SETS

### BASIC DEFINITIONS AND FACTS

**Cardinality definition** Sets  $A$  and  $B$  have the same cardinality iff  $\exists f( f : A \xrightarrow{1-1, onto} B)$ .

**Cardinality notations** If sets  $A$  and  $B$  have the same cardinality we denote it as:  $|A| = |B|$  or  $cardA = cardB$ , or  $A \sim B$ . We also say that  $A$  and  $B$  are *equipotent*.

**Cardinality** We put the above notations together in one definition:

$|A| = |B|$  or  $cardA = cardB$ , or  $A \sim B$  iff  $\exists f( f : A \xrightarrow{1-1, onto} B)$ .

**Finite** A set  $A$  is finite iff  $\exists n \in \mathbb{N} \exists f( f : \{0, 1, 2, \dots, n-1\} \xrightarrow{1-1, onto} A)$ , i.e. we say: a set  $A$  is finite iff  $\exists n \in \mathbb{N}(|A| = n)$ .

**Infinite** A set  $A$  is infinite iff  $A$  is NOT finite.

**Aleph zero**  $\aleph_0$  (Aleph zero) is a cardinality of  $\mathbb{N}$  (Natural numbers).

**For any set**, set  $A$  has a cardinality  $\aleph_0$  ( $|A| = \aleph_0$ ) iff  $A \sim \mathbb{N}$ , (or  $|A| = |\mathbb{N}|$ , or  $cardA = card\mathbb{N}$ ).

**Countable** A set  $A$  is countable iff  $A$  is finite or  $|A| = \aleph_0$ .

**Infinitely countable** A set  $A$  is infinitely countable iff  $|A| = \aleph_0$ .

**Uncountable** A set  $A$  is uncountable iff  $A$  is NOT countable.

**Observe** that it means that

A set  $A$  is uncountable iff  $A$  is infinite and  $|A| \neq \aleph_0$ .

**Continuum**  $\mathcal{C}$  (Continuum) is a cardinality of Real numbers, i.e.  $\mathcal{C} = |\mathbb{R}|$ .

**We sat that** a set  $A$  has a cardinality  $\mathcal{C}$  ( $|A| = \mathcal{C}$ ) iff  $|A| = |\mathbb{R}|$ .

**Cardinality  $A \leq$  cardinality  $B$**  We define  $|A| \leq |B|$  iff  $A \sim C$  and  $C \subseteq B$ .

**Simple Fact** If  $A \subseteq B$  then  $|A| \leq |B|$ .

**For any cardinal numbers  $\mathcal{N}, \mathcal{M}$** , we say that

$\mathcal{N} \leq \mathcal{M}$  iff for any sets  $A, B$ , such that  $|A| = \mathcal{N}$  and  $|B| = \mathcal{M}$  we have  $|A| \leq |B|$ .

**Cardinality  $A <$  cardinality  $B$**   $|A| < |B|$  iff  $|A| \leq |B|$  and  $|A| \neq |B|$ .

**For any cardinal numbers  $\mathcal{N}, \mathcal{M}$**  we say that

$\mathcal{N} < \mathcal{M}$  iff for any sets  $A, B$ , such that  $|A| = \mathcal{N}$  and  $|B| = \mathcal{M}$  we have  $|A| < |B|$ .

**Cantor Theorem** For any set  $A$ ,  $|A| < |\mathcal{P}(A)|$ .

**Cantor-Berstein Theorem** For any sets  $A, B$ ,

If  $|A| \leq |B|$  and  $|B| \leq |A|$ , then  $|A| = |B|$ .

For any cardinal numbers  $\mathcal{N}, \mathcal{M}$ , we have that

If  $\mathcal{N} \leq \mathcal{M}$  and  $\mathcal{M} \leq \mathcal{N}$ , then  $\mathcal{N} = \mathcal{M}$ .

### ARITHMETIC OF CARDINAL NUMBERS

**Sum** ( $\mathcal{N} + \mathcal{M}$ ) We define:

$\mathcal{N} + \mathcal{M} = |A \cup B|$ , where  $A, B$  are such that  $|A| = \mathcal{N}$ ,  $|B| = \mathcal{M}$  and  $A \cap B = \emptyset$ .

**Multiplication** ( $\mathcal{N} \cdot \mathcal{M}$ ) We define:

$\mathcal{N} \cdot \mathcal{M} = |A \times B|$ , where  $A, B$  are such that  $|A| = \mathcal{N}$ ,  $|B| = \mathcal{M}$ .

**Power** ( $\mathcal{M}^{\mathcal{N}}$ )  $\mathcal{M}^{\mathcal{N}} = \text{card}\{f : f : A \rightarrow B\}$ , where  $A, B$  are such that  $|A| = \mathcal{N}$ ,  $|B| = \mathcal{M}$ .

**Observe** that the definition says that  $\mathcal{M}^{\mathcal{N}}$  is the cardinality of all functions that map a set  $A$  (of cardinality  $\mathcal{N}$ ) into a set  $B$  (of cardinality  $\mathcal{M}$ ).

**Power**  $2^{\mathcal{N}}$  We define:

$2^{\mathcal{N}} = \text{card}\{f : f : A \rightarrow \{0, 1\}\}$ , where  $|A| = \mathcal{N}$ .

$2^{\mathcal{N}}$  **Theorems** We prove the following.

1.  $2^{\mathcal{N}} = \text{card}\mathcal{P}(A)$ , where  $|A| = \mathcal{N}$ .

2.  $2^{\aleph_0} = \mathcal{C}$ .

**Power Properties**  $\mathcal{N}^{\mathcal{P}+\mathcal{T}} = \mathcal{N}^{\mathcal{P}} \cdot \mathcal{N}^{\mathcal{T}}$ .  $(\mathcal{N}^{\mathcal{P}})^{\mathcal{T}} = \mathcal{N}^{\mathcal{P} \cdot \mathcal{T}}$ .

### ARITHMETIC OF $n, \aleph_0, \mathcal{C}$

**Union 1**  $\aleph_0 + \aleph_0 = \aleph_0$ .

Union of two infinitely countable sets is an infinitely countable set.

**Union 2**  $\aleph_0 + n = \aleph_0$ .

Union of a finite (cardinality  $n$ ) and infinitely countable set is an infinitely countable set.

**Union 3**  $\aleph_0 + \mathcal{C} = \mathcal{C}$ .

Union of an infinitely countable set and a set of the same cardinality as Real numbers has the same cardinality as the set of Real numbers.

**Union 4**  $\mathcal{C} + \mathcal{C} = \mathcal{C}$ .

Union of two sets of cardinality the same as Real numbers has the same cardinality as the set of Real numbers.

**Cartesian Product 1**  $\aleph_0 \cdot \aleph_0 = \aleph_0$ .

Cartesian Product of two infinitely countable sets is an infinitely countable set.

**Cartesian Product 2**  $n \cdot \aleph_0 = \aleph_0$ .

Cartesian Product of a finite set and an infinitely countable set is an infinitely countable set.

**Cartesian Product 3**  $\aleph_0 \cdot \mathcal{C} = \mathcal{C}$ .

Cartesian Product of an infinitely countable set and a set of the same cardinality as Real numbers has the same cardinality as the set of Real numbers.

**Cartesian Product 4**  $\mathcal{C} \cdot \mathcal{C} = \mathcal{C}$ .

Cartesian Product of two sets of cardinality the same as Real numbers has the same cardinality as the set of Real numbers.

**Power 1**  $2^{\aleph_0} = \mathcal{C}$ .

The set of all subsets of Natural numbers (or any set equipotent with natural numbers) has the same cardinality as the set of Real numbers.

**Power 2**  $\aleph_0^{\aleph_0} = \mathcal{C}$ .

There are  $\mathcal{C}$  of all functions that map  $\mathbb{N}$  into  $\mathbb{N}$ .

There are  $\mathcal{C}$  sequences (all sequences) that can be form out of an infinitely countable set.

$$\aleph_0^{\aleph_0} = \{f : f : \mathbb{N} \longrightarrow \mathbb{N}\} = \mathcal{C}.$$

**Power 3**  $\mathcal{C}^{\mathcal{C}} = 2^{\mathcal{C}}$ .

There are  $2^{\mathcal{C}}$  of all functions that map  $\mathbb{R}$  into  $\mathbb{R}$ .

The set of all real functions of one variable has the same cardinality as the set of all subsets of Real numbers.

**Inequalities**  $n < \aleph_0 < \mathcal{C}$ .

**Theorem** If  $A$  is a finite set,  $A^*$  is the set of all finite sequences formed out of  $A$ , then  $A^*$  has  $\aleph_0$  elements.

Shortly: If  $|A| = n$ , then  $|A^*| = \aleph_0$ .