

Conformal Spherical Parametrization for High Genus Surfaces

Wei Zeng*
Chinese Academy of Sciences
Stony Brook University

Xin Li†
Stony Brook University

Shing-Tung Yau‡
Harvard University

Xianfeng Gu§
Stony Brook University

Abstract

Surface parameterization establishes bijective maps from a surface onto a topologically equivalent standard domain. It is well known that the spherical parameterization is limited to genus-zero surfaces. In this work, we design a new parameter domain, two-layered sphere, and present a framework for mapping high genus surfaces onto sphere. This setup allows us to transfer the existing applications based on general spherical parameterization to the field of high genus surfaces, such as remeshing, consistent parameterization, shape analysis, and so on.

Our method is based on Riemann surface theory. We construct meromorphic functions on surfaces: for genus one surfaces, we apply Weierstrass P-functions; for high genus surfaces, we compute the quotient between two holomorphic one-forms.

Our method of spherical parameterization is theoretically sound and practically efficient. It makes the subsequent applications on high genus surfaces very promising.

CR Categories: I.3.5 [Computer Graphics]: Computational Geometry and Object Modeling—Curve, Surface and Solid and Object Representations.

Keywords: Conformal Spherical Parameterization, Meromorphic Function, High Genus Surface, Layered Sphere

1 Introduction

Surface parameterization (for a recent survey, we refer the reader to [Floater and Hormann 2005]) is a fundamental tool in computer graphics and benefits many digital geometry processing applications such as texture mapping, shape analysis, compression, morphing, remeshing, etc. Some problems become much easier to deal with a uniform parameter domain. Usually in these settings surfaces are represented as triangle meshes, and the maps are required to be at least no-foldovers and low-distortion in terms of area, angle, or both aspects.

In graphics, spherical parameterizations for genus zero closed surfaces have been proposed and widely used in the past. Most methods [Gotsman et al. 2003; Gu et al. 2004; Haker et al. 2000; Sheffer et al. 2004; Praun and Hoppe 2003] are to directly map the mesh to spherical domain, which is usually formulated as a spherical energy minimization problem, such as conformal, Tutte, Dirichlet, area, spring, stretch energies, or their combinations, as cited in [Floater and Hormann 2005]. The optimization process is to relax the initial map to reach no-foldovers under specified distortion metric.

In medical imaging, spherical parameterizations are broadly applied for brain cortex surface mapping. In this setting, preservation of local shapes are crucial. Therefore, different conformal spherical parameterizations are proposed. Angenent et.al. [Angenent et al. 1999] construct meromorphic functions on the brain surface directly, then lift the mapping onto the sphere using inverse stereographic projections. Gu et.al. [Gu et al. 2004] compute harmonic maps between the brain cortex surface and the unit sphere and use Möbius transformation to adjust the map. Stephenson [Stephenson 2005] uses circle packing method to construct conformal brain mapping.

However, it is well known that the spherical parameterization is limited to genus-zero models. To the best of our knowledge, there are few works on high genus surfaces. Recently, Lee et.al. [Lee et al. 2006] present a construction method by Boolean operations of positive and negative spheres. This method requires a lot of interactive human recognitions and geometry editing techniques. Furthermore, the results are not conformal.

In this work, we aim at automatic generalizing conformal spherical parameterizations for high genus surfaces. Because high genus surfaces and spheres are not topologically equivalent, we allow the existence of branch points.

Our method relies on the conformal structure for higher genus meshes. There are two ways to compute conformal structures of general surfaces: one method is based on Hodge theory [Gu and Yau 2002], and the other on discrete surface Ricci flow [Gu et al. 2005; Jin et al. 2006a; Jin et al. 2006b].

According to Riemann surface theory, a conformal map between a surface and the sphere is equivalent to a meromorphic function defined on the surface. The map wraps the surface onto the sphere by several layers and has several branch points, the number of layers and the branch points are determined by the topology of the surface (by Riemann-Hurwitz theorem). The key is how to construct the meromorphic functions on the input surface. For genus one closed surfaces, we construct the well-known Weierstrass P-function. For high genus surfaces, the quotient between two holomorphic 1-forms is a meromorphic function.

Compared with the existing planar parameterization for high genus meshes, the layered sphere is more natural domain than the planar domain. Employing the properties of sphere geometry and the existing spherical parameterization related applications on genus-zero meshes, the

*e-mail: zengwei@cs.sunysb.edu: Institute of Computing Technology, Chinese Academy of Sciences; Graduate school of the Chinese Academy of Sciences

†e-mail: xinli@cs.sunysb.edu

‡e-mail: yau@math.harvard.edu

§e-mail: gu@cs.sunysb.edu

spherical parameterization designed for high genus meshes can get more insights on shape analysis, and introduce more possible applications for high genus meshes.

The contributions of this work are briefly as follows:

- To present a novel practical framework to compute conformal spherical parameterizations for general surfaces;
- To extend the applications of general spherical parameterization onto that of high genus meshes, including re-meshing, morphing, etc;
- To introduce a systematic method to compute meromorphic functions on general Riemann surfaces.

The remainder of the paper is organized as follows: Section 2 illustrates the basic definition and theorem in theory used in this work, Section 3 describes the algorithm flow, and Section 4 shows the experimental results, and discussion on the method. The paper concludes with a conclusion and future work in Section 5.

2 Basic Theory

In this section, we will briefly introduce the basic theories of meromorphic function on Riemann surfaces. We refer readers to [Farkas and Kra 1980] for details.

2.1 Riemann Surfaces

Definition 2.1. *Manifold.* Suppose M is a topological space, each point has a neighborhood U_α and a *homeomorphism* $\phi_\alpha : U_\alpha \rightarrow V_\alpha$ from U_α to an open set V_α in \mathbf{R}^n . (U_α, ϕ_α) is called a local chart. If two such neighborhoods U_α, U_β intersect, then the chart transition function

$$\phi_{\alpha\beta} = \phi_\beta \phi_\alpha^{-1} : \phi_\alpha(U_\alpha \cap U_\beta) \rightarrow \phi_\beta(U_\beta \cap U_\alpha)$$

is a homeomorphism from one open set of \mathbf{R}^n to another. Then M is a n dimensional manifold, the set of all local charts $\{(U_\alpha, \phi_\alpha)\}$ form an atlas.

Definition 2.2. *Holomorphic functions.* In complex analysis, a function $f : \mathbb{C} \rightarrow \mathbb{C}$, $(x, y) \rightarrow (u, v)$ is *holomorphic*, if and only if it satisfies the following Riemann-Cauchy equation

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}.$$

Definition 2.3. *Riemann Surfaces.* A Riemann surface is a two dimensional manifold with an atlas $\{(U_\alpha, \phi_\alpha)\}$, such that all chart transitions $\phi_{\alpha\beta}$ are holomorphic functions. The atlas is called the conformal atlas, and each local coordinates $\phi_\alpha(U_\alpha)$ are called holomorphic coordinates. The maximal conformal atlas is called a conformal structure of the surface.

Definition 2.4. *Riemannian Metric.* A Riemannian metric on a manifold is a tensor \mathbf{g} , which defines an inner product $\langle \cdot, \cdot \rangle_{\mathbf{g}}$ on the tangent spaces of the manifold. Suppose v_1, v_2 are two tangent vectors on the tangent space at point p , then the angle between them can be calculated as

$$\theta = \cos^{-1} \frac{\langle v_1, v_2 \rangle_{\mathbf{g}}}{\sqrt{\langle v_1, v_1 \rangle_{\mathbf{g}}} \sqrt{\langle v_2, v_2 \rangle_{\mathbf{g}}}}.$$

In the current work, we focus on surfaces embedded in the Euclidean space \mathbb{R}^3 . Therefore, all the surfaces are with induced Euclidean metric \mathbf{g} . We require the conformal structure and the Riemannian metric be compatible in the following way. Suppose $\{(U_\alpha, \phi_\alpha)\}$ is a conformal atlas, we denote the local coordinates of (U_α, ϕ_α) as (x_α, y_α) , then the metric tensor is represented

$$ds^2 = e^{2\lambda(x_\alpha, y_\alpha)}(dx_\alpha^2 + dy_\alpha^2),$$

where $\lambda(x_\alpha, y_\alpha)$ is a function. It can be easily verified that the intersection angle measured by \mathbf{g} equals to that measured by Euclidean metric defined on the parameter domain

$$d\bar{s}^2 = dx_\alpha^2 + dy_\alpha^2.$$

Namely, the local coordinates preserve angles; therefore conformal structure is also called angle-preserving structure.

2.2 Holomorphic 1-forms

Our algorithm heavily depends on the calculation of holomorphic 1-forms of a given Riemann surface.

Definition 2.5. *Holomorphic 1-forms.* Given a Riemann surface X with a conformal atlas (U_α, z_α) , a *holomorphic 1-form* ω is defined by a family $(U_\alpha, z_\alpha, \omega_\alpha)$, such that (1) $\omega_\alpha = f_\alpha(z_\alpha)dz_\alpha$, where f_α is holomorphic on U_α , and (2) if $z_\alpha = \phi_{\alpha\beta}(z_\beta)$ is the coordinate transformation on $U_\alpha \cap U_\beta (\neq \emptyset)$, $f_\alpha(z_\alpha) \frac{dz_\alpha}{dz_\beta} = f_\beta(z_\beta)$, the local representation of the differential form ω satisfies the chain rule.

For a Riemann surface X with genus- $g (g > 0)$, all holomorphic 1-forms on X form a complex g -dimensional vector space ($2g$ real dimensions), denoted as $\Omega^1(X)$. The quality of a global conformal parameterization for a high genus surface is mainly determined by the choice




Figure 1: Example of holomorphic 1-forms. For genus-two cases, there are four 1-forms, described by the check-board texture mapping. The parameter (u, v) of each vertex is the average of 1-forms of its connected edges.

of the holomorphic 1-form. The zero points of a holomorphic 1-form ω are the points where, on any local representation $(U_\alpha, z_\alpha, \omega_\alpha)$, ω_α equals zero. For a genus- g ($g > 0$) surface, there are in general $2g - 2$ zero points for each holomorphic 1-form. Figure 1 illustrates the holomorphic 1-forms on a genus-two surface.

2.3 Meromorphic Functions

Definition 2.6. Holomorphic Map. Suppose X, Y are Riemann surfaces with conformal atlases $\{(U_\alpha, \phi_\alpha)\}$ and $\{(V_\beta, \psi_\beta)\}$, a holomorphic map between X and Y is a continuous map $f : X \rightarrow Y$ such that for each holomorphic coordinates ϕ_α on U_α containing x on X and ψ_β defined in a neighborhood of $f(x)$ on Y , the composition

$$\psi_\beta \circ f \circ \phi_\alpha^{-1}$$

is holomorphic.

Intuitively, a holomorphic map is a conformal (angle preserving) map between two surfaces. In general, such map doesn't exist between two high genus surfaces. But conformal maps between a high genus to the unit sphere (with branch points) always exist.

Definition 2.7. Meromorphic Function. A meromorphic function f on a Riemann surface X is a holomorphic map to the Riemann sphere $S = \mathbb{C} \cup \{\infty\}$.

The meromorphic function on a Riemann surface usually has multiple branch points. A branch point may be informally thought of as a point at which a "multiple-valued function" changes values when one winds once around it. The neighborhood of branch points wrap around the range a finite or infinite number of times.

Definition 2.8. Branch Point. A branch point of an analytic function is a point in the complex plane whose complex argument can be mapped from a single point in the domain to multiple points in the range.

For example, if $n > 1$, then z^n has a degree n branch point at 0.

The relationship between the Euler characteristics of the source surface and the target surface is described by the following theorem, when one is a covering of the other (here, using Riemann sphere) with branch points.

Theorem 2.9. Riemann-Hurwitz. Let $f : X \rightarrow S$ be a meromorphic function of degree d on a closed connected Riemann surface X , and suppose it has branch points x_1, \dots, x_n where the local form of $f(x) - f(x_k)$ is a holomorphic function with a zero of multiplicity m_k . Then

$$\chi(X) = 2d - \sum_{k=1}^n (m_k - 1)$$

where $\chi(X)$ is the Euler number of X and there is $\chi(S) = 2$.

In the settings of this paper, the function is two-valued, there are two overlapped layers on the complex plane, which is lifted to be a *two-layered sphere* by inverse stereo graphic projection, seen in Figure 2. Thus, from the Riemann-Hurwitz formula, the number of branch points can be computed indirectly. Given a closed g -holed tori, there are $2g + 2$ branch points (See Figure 3, 4).

3 Algorithm

This section explains the algorithm for computing the conformal spherical parameterizations for high genus surfaces based on meromorphic functions.

3.1 Holomorphic 1-forms

At the first stage, we calculate the basis for the holomorphic 1-form group $\Omega^1(M)$. The method is based on the Hodge theory. All the surfaces are represented as triangular meshes (simplicial complex). The followings are the major steps, for details we refer readers to [Gu and Yau 2003]. We assume the input surface is a genus g closed surface.




Figure 2: Two-layered Sphere for genus-one and genus-two cases of our method. From left to right, they are (1) one-hole torus mesh with 10,000 vertices and 20,000 faces, (2) two-layered sphere with four branch points, (3) two-holes torus mesh with 12,286 vertices and 24,576 faces, and (4) two-layered sphere with six branch points. Two layers are connected by branch points where the lines twist together.




Figure 3: Branch points illustration for genus- g ($g > 0$) surfaces. There are $2g + 2$ branch points, two for each tunnel. The outer space is regarded as a tunnel here.

1. Compute the basis of the first homology group $H_1(M, \mathbb{Z})$, $\{\gamma_1, \gamma_2, \dots, \gamma_{2g}\}$.
2. Compute the dual cohomology group basis $H^1(M, \mathbb{Z})$, $\{\omega_1, \omega_2, \dots, \omega_{2g}\}$, such that $\omega_i(\gamma_j) = \delta_j^i$.
3. Diffuse cohomology basis to harmonic 1-forms, such that $\Delta\omega_i = 0$, where Δ is the Laplace-Beltrami operator.
4. Compute the conjugate harmonic 1-forms, ω_i^* , where $*$ is the Hodge star operator. $\{\omega_1 + \sqrt{-1}\omega_1^*, \omega_2 + \sqrt{-1}\omega_2^*, \dots, \omega_{2g} + \sqrt{-1}\omega_{2g}^*\}$ form a basis for the holomorphic 1-form group.

3.2 Genus One Surfaces

For a genus one closed surface M , we first compute the holomorphic 1-form basis, we denote a holomorphic 1-form as ω . ω induces a Riemannian metric

$$ds^2 = \omega\bar{\omega},$$

which is flat everywhere, namely, the Gaussian curvature induced by ω is zero.

Definition 3.1. Universal Covering Space. Suppose M is a surface, $\pi: \bar{M} \rightarrow M$ is a covering space if for every point p in M there is a neighborhood U of p so that $\pi^{-1}(U)$ is a disjoint union of open sets V_i , $i \in I$, and the restriction $\pi|_{V_i}: V_i \rightarrow U$ is a homeomorphism for each i . If \bar{M} is simply connected, then $\pi: \bar{M} \rightarrow M$ is a universal covering space.

Definition 3.2. Deck Transformation. Suppose M is a surface, the pair (\bar{M}, π) is the universal covering space of M , $\phi: \bar{M} \rightarrow \bar{M}$ is called a deck transformation, if $\pi \circ \phi = \pi$.

All the deck transformations of a surface form a group. For the one-hole torus case, we can embed the universal covering space onto the complex plane using the flat metric. Then the deck transformations of (\bar{M}, ds^2) are translations. We compute the generators of the deck transformation group $Deck(M)$, which are translations, and denote them as $\{w_1, w_2\}$. Any deck transformation can be represented as $w = mw_1 + nw_2, m, n \in \mathbb{Z}$.

The meromorphic function can be easily constructed using the Weierstrass P-function on the planar domain. Define

$$\wp(z) = \frac{1}{z^2} + \sum_{w \neq 0} \left(\frac{1}{(z-w)^2} - \frac{1}{w^2} \right)$$

where the sum is over all non-zero $w \in Deck(M)$. Because the sum is essentially over all deck transformations, therefore the Weierstrass function is invariant under the action of $Deck(M)$,

$$\wp(z+w) = \wp(z), \forall w \in Deck(M),$$

so that this is a meromorphic function defined on the surface M . The branch points of $\wp(z)$ are $0, \frac{w_1}{2}, \frac{w_2}{2}, \frac{w_1+w_2}{2}$, (see Figure 5 and 6). The computation example can be seen in Figure 7.




Figure 4: Branch points of torus. The branch points are shown by the texture, where the region is formed by eight edges. The corresponding location is shown in its right by practical method.




Figure 5: Generator (w_1, w_2) of deck transformations. The branch points here are labeled by four solid circle points, denoting $0, \frac{w_1}{2}, \frac{w_2}{2},$ and $\frac{w_1+w_2}{2}$ respectively.

3.3 Genus- $g(g > 1)$ Surfaces

Suppose M is a genus g closed surface, $\{\omega_1, \omega_2, \dots, \omega_{2g}\}$ form a basis of the holomorphic 1-form group. A meromorphic function can be constructed using the ratio between two holomorphic 1-forms.

Suppose τ_1 and τ_2 are two holomorphic 1-forms,

$$\tau_k = \sum_j a_{kj} \omega_j, k = 1, 2.$$

Suppose $\{(U_\alpha, \phi_\alpha)\}$ is the conformal atlas of the surface M , on a local chart (U_α, ϕ_α) ,

$$\tau_1 = f_1(z_\alpha) dz_\alpha, \tau_2 = f_2(z_\alpha) dz_\alpha, \quad (1)$$

the ratio is

$$F(z_\alpha) = \frac{f_1(z_\alpha)}{f_2(z_\alpha)}.$$

On an overlapping chart (U_β, ϕ_β) , it can be easily verified that $F(z_\beta) = F(z_\alpha(z_\beta))$, therefore F is a globally well defined meromorphic function on M .

We construct the meromorphic function as $F = \frac{\omega_1 + \omega_2}{\omega_1 - \omega_2}$. The computation example can be seen in Figure 8.

3.4 Branch Point Location

According to Riemann-Hurwitz theorem, meromorphic function on a Riemann surface has branch points, which are important geometric features of the conformal structure of the surface. In practice, it is crucial to accurately locate these branch points.

In order to locate the branch points, we apply the following algorithm to compute the winding number of a closed planar curve. For each point $p \in M$ on the surface, $F(p) \in \mathbb{C}$ is on the complex plane. We choose a simple closed curve γ surrounding p on M , then $F(\gamma)$ is a curve on \mathbb{C} , we define the winding number as

$$I(\gamma, p) = \frac{1}{2\pi\sqrt{-1}} \int_{F(\gamma)} \frac{dz}{z - F(p)}.$$

If $I(\gamma, p) \neq \pm 1$, then p is a branch point.

In practice, in order to improve the accuracy of computing the winding number, we can zoom the neighborhood of a point p by a Möbius transformation,

$$\phi : \mathbb{C} \rightarrow \mathbb{C}, \phi(z) = \frac{az + b}{cz + d}, a, b, c, d \in \mathbb{C}, ad - bc = 1.0.$$




Figure 6: Universal covering space and Weierstrass P-function. From left to right, they are (1) kitten mesh, (2) part of universal covering space, including 24 parameter periods, (3) one parameter period, and (4) parameter domain under Weierstrass p-function.




Figure 7: Genus-one case by Weierstrass p-function. For one-hole torus case, the computation steps are (from left to right) (1) one parameter period with zoomed-in part (right), (2) parameter domain under Weierstrass p-function with zoomed-in part (right), (3) sphere by inverse stereographic projection, (4) parameter domain after Möbius transformation, and (5) sphere of (4) by inverse stereographic projection.

In order to map the complex plane to the sphere conformally, we use the stereographic projection, suppose a point (x, y, z) is on the unit sphere \mathbb{S}^2 . It is mapped to the planar point $(\frac{2x}{z-1} + \sqrt{-1} \frac{2y}{z-1})$.

In practice, for different purposes, in order to improve the quality of the spherical conformal surface parameterizations, we can choose the holomorphic 1-forms τ_1, τ_2 in Equation 1, and use Möbius transformation to further improve the parameterizations.

4 Results and Discussion

In our implementation, all surfaces are triangular meshes and represented as half edge data structure. The holomorphic 1-forms are represented as simplicial co-chains on the mesh. The whole algorithm is implemented using c++ on windows platform.

Under the theoretical support of meromorphic function, the implementation of our method needs only a simple extension of existing conformal planar mesh parametrization methods

[Gu and Yau 2002; Gu and Yau 2003]. The results are bijective maps (except at the branch points) and conformal as illustrated in figures 9, 10, 11, and 12, with colored check-board texture.

We conducted experiments on a variety of meshes. Since our method is based on a composition of several maps, planar parameterization, stereographic projection and conformal relaxation, the validity and conformality of each them guarantees the validity and conformality of the whole map. Using the valid embedding and the periodicity of the planar parameterization, the validity is guaranteed except the overlapping around the branch points.

For high genus surfaces, the branch points can be detected technically or computed from theoretical definition. Their number b is determined by the number h of tunnels. Here, we regard the outer boundary as an outer tunnel. So, for closed two-manifolds, there are formula $h = g + 1$, and $b = 2h$. You can easily get this view from that there are two branch points for each tunnel boundary. Figure 4 illustrates the experimental results for genus-one, genus-two surfaces. For genus-one case, their exact locations depend on the start point of computation, because of the periodicity of parameterization domain. Different choices of ω_1 and ω_2 get different branch points.

From the planar illustration (Figure 3), you can get the insight that the whole surface is split into two parts by the sequence connection of each branch points, labeled in dashed lines. Each part is corresponding to a layer on the spherical domain, which can be seen in Figure 9, where the parameter (u, v) is the spherical curvilinear coordinate (θ, ϕ) [Zayer et al. 2006].

As the most direct application of parameterization, texture mapping results are shown in Figure 9, 10, and 11. The conformal parameters (u, v) by stereographic projection may cover a lot of periods of the texture image, so there are many duplicated texture periods.

Compared with Gaussian spherical mapping, our method embeds the high genus surfaces onto sphere with one twisting layer (or multiple layers connecting with branch points) and shows one more natural way for their spherical representations. The simple Gaussian mapping generates the overlapping on handles and can not be embedded properly even under powerfully iterative harmonic relaxation. The handles finally shrink to dense bands, see Figure 13, i.e., the handle information is lost, where under-sampling occurs.




Figure 8: Genus- n ($n > 1$) case by holomorphic 1-forms. For two-holes torus case, the computation steps are (from left to right) (1) two holomorphic 1-forms ω_1 and ω_2 , (2) parameter domain under quotient function by ω_1 and ω_2 , (3) sphere by inverse stereographic projection, (4) parameter domain after Möbius transformation, and (5) sphere of (4) by inverse stereographic projection.




Figure 9: Spherical conformal parameterization for one-hole torus case. The 1st and 2nd rows are mapped by different textures. From left and to right, each column denotes: (1) conformal parameterization result on layered sphere, (2) initial spherical conformal parameterization result on original surface, (3) spherical conformal parameterization result with Möbius transformation, and (4) curvilinear parameterization of (1), and (5) curvilinear parameterization of (2).

5 Conclusions and Future Work

Conventional conformal spherical parameterizations are limited to genus-zero surfaces. In this work, we generalized conformal spherical parameterizations to surfaces with arbitrary topologies. The proposed method is based on constructing meromorphic functions on Riemann surfaces. High genus surfaces are conformally mapped to the unit sphere with branching points, which are the key features for the conformal structure of the surface. The method is theoretically sound and practically efficient.

In the future, we will explore along the following directions:

- Intrinsic triangulations. Triangulate the spherical images with branch points, such that all branch points are vertices, pull back the triangulation to the original surface. This method gives us an intrinsic triangulation, which is solely determined by the conformal structure of the surface.
- Surface matching based on the consistent spherical triangulation. If two surfaces are with similar conformal structures, they can be triangulated in a consistent way using the above intrinsic triangulations. The related applications around consistent parameterization [Praun et al. 2001] can be developed onto high genus surfaces, such as morphing, mapping, matching, etc.
- Meromorphic function based on Poincaré series. Current meromorphic functions on high genus surfaces are the quotient of holomorphic 1-forms, we will develop novel method based on Poincaré series.

References

- ANGENENT, S., HAKER, S., TANNENBAUM, A., AND KIKINIS, R. 1999. Conformal geometry and brain flattening. In *MICCAI '99: Proceedings of the Second International Conference on Medical Image Computing and Computer-Assisted Intervention*, Springer-Verlag, London, UK, 271–278.
- ASIRVATHAM, A., PRAUN, E., AND HOPPE, H. 2005. Consistent spherical parameterization. In *International Conference on Computational Science (2)*, 265–272.




Figure 10: Spherical conformal parameterization for two-holes torus case. The 1st and 2nd rows are mapped by different textures. From left to right, each column denotes: (1) conformal parameterization result on layered sphere, (2) initial spherical conformal parameterization result on original surface, (3) spherical conformal parameterization result with Möbius transformation, and




Figure 11: Spherical conformal parameterization for kitten case with 10,219 vertices and 20,438 faces. The 1st and 2nd rows illustrate both sides of the model. From left to right, each column denotes: (1) and (2) are initial spherical conformal parameterization results on original surface; (3) and (4) are spherical conformal parameterization results with Möbius transformation; (5) shows the branch points.

FARKAS, H. M., AND KRA, I. 1980. *Riemann Surfaces*. Springer-Verlag, New York.

FLOATER, M. S., AND HORMANN, K., 2005. Surface parameterization: a tutorial and survey.

GOTSMAN, C., GU, X., AND SHEFFER, A. 2003. Fundamentals of spherical parameterization for 3d meshes. *ACM Trans. Graph.* 22, 3, 358–363.

GU, X., AND YAU, S. 2002. Computing conformal structures of surfaces. In *Communications in Information and Systems*, vol. 2, 121–146.

GU, X., AND YAU, S. 2003. Global conformal surface parameterization. In *SGP '03: Proceedings of the 2003 Eurographics/ACM SIGGRAPH symposium on Geometry processing*, Eurographics Association, Aire-la-Ville, Switzerland, Switzerland, 127–137.

GU, X., WANG, Y., CHAN, T. F., THOMPSON, P. M., AND YAU, S.-T. 2004. Genus zero surface conformal mapping and its application to brain surface mapping. *IEEE Transaction on Medical Imaging* 23, 8, 949–958.

GU, X., LUO, F., AND YAU, S.-T. 2005. Tutorial on discrete ricci flow for global parameterizations. Technique report, August.

HAKER, S., ANGENENT, S., TANNENBAUM, A., KIKINIS, R., SAPIRO, G., AND HALLE, M. 2000. Conformal surface parameterization for texture mapping. *IEEE Transactions on Visualization and Computer Graphics* 6, 2, 181–189.

JIN, M., KIM, J., LUO, F., LEE, S., AND GU, X. 2006. Conformal surface parameterization using euclidean ricci flow. Technique report, May.




Figure 12: Spherical conformal parameterization for vase1 and vase2 cases, with the same number vertices and faces. From left to right, each column is (1) triangular meshes with 5,910 vertices and 11,824 faces; (2) initial spherical conformal parameterization results on original surface; (3) the results with Möbius transformation.




Figure 13: Gaussian mapping under harmonic optimization with iterative times 0, 500, and 1200.

- JIN, M., LUO, F., AND GU, X. 2006. Computing surface hyperbolic structure and real projective structure. In *SPM '06: Proceedings of the 2006 ACM symposium on Solid and physical modeling*, ACM Press, New York, NY, USA, 105–116.
- KRAEVOY, V., AND SHEFFER, A. 2004. Cross-parameterization and compatible remeshing of 3d models. *ACM Trans. Graph.* 23, 3, 861–869.
- LEE, T.-Y., YAO, C.-Y., CHU, H.-K., TAI, M.-J., AND CHEN, C.-C. 2006. Generating genus- n -to- m mesh morphing using spherical parameterization. *Computer Animation and Virtual Worlds* 17, 3-4, 433–443.
- PRAUN, E., AND HOPPE, H. 2003. Spherical parametrization and remeshing. In *SIGGRAPH '03: ACM SIGGRAPH 2003 Papers*, ACM Press, New York, NY, USA, 340–349.
- PRAUN, E., SWELDENS, W., AND SCHRÖDER, P. 2001. Consistent mesh parameterizations. In *SIGGRAPH 2001, Computer Graphics Proceedings*, ACM Press / ACM SIGGRAPH, E. Fiume, Ed., 179–184.
- SHEFFER, A., GOTSMAN, C., AND DYN, N. 2004. Robust spherical parameterization of triangular meshes. *Computing* 72, 1-2, 185–193.
- STEPHENSON, K. 2005. *Introduction to Circle Packing: the Theory of Discrete Analytic Functions*. Camb. Univ. Press, New York. ISBN 0-521-82356-0, QA640.7.S74.
- ZAYER, R., RÖSSL, C., AND SEIDEL, H.-P. 2006. Curvilinear spherical parameterization. In *Shape Modeling International (SMI)*, IEEE, Matsushima, Japan, 57–64.