

CSE303 Q1 SOLUTIONS

Quiz has TWO PARTS

Part 1: yes/no questions (15pts)

Must write justifications- **no justification, no points!**

Part 2 (10pts) Solve only one problem of your choice

PART 1: Yes/No Questions

- $\{\emptyset\} \subseteq \{a, b, c\}$
Justify: $\emptyset \notin \{a, b, c\}$ **n**
- $2^{\{1,2\}} \cap \{\{1,2\}\} \neq \emptyset$
Justify: $\{1,2\} \subseteq \{1,2\}$ i.e. $\{1,2\} \in 2^{\{1,2\}}$ and $\{1,2\} \in \{\{1,2\}\}$ **y**
- Some $R \subseteq A \times B$ are functions that map A into B
Justify: Functions are special type of relations. **y**
- Let $A \neq \emptyset$ such that there are exactly 25 partitions of A . It is possible to define 20 equivalence relations on A
Justify: one can define up to 25 (as many as partitions) of equivalence classes **y**
- There is a relation that is equivalence and *order* at the same time
Justify: equality relation **y**
- Let $A = \{n \in \mathbb{N} : n^2 + 1 \leq 15\}$. It is possible to define 8 alphabets $\Sigma \subseteq A$
Justify: A has 4 elements, so we have $2^4 > 8$ subsets **y**
- Let $A = \{\{n\} \in 2^{\mathbb{N}} : n^2 + 1 \leq 15\}$. A is infinite.
Justify: $\{n \in \mathbb{N} : n^2 + 1 \leq 15\} = \{0, 1, 2, 3\}$, hence $A = \{\{0\}, \{1\}, \{2\}, \{3\}\}$ is a finite set **n**
- Let $\Sigma = \{a\}$. There are countably many languages over Σ .
Justify: There is any many languages as subsets of Σ^* , i.e. uncountably many and exactly as many as real numbers. **n**
- Let $\Sigma = \{n \in \mathbb{N} : n \leq 1\}$
There are uncountably many finite languages over Σ
Justify: $\Sigma = \{0, 1\}$ and so Σ^* is countably infinite. The set of all finite subsets of any countably infinite set is countably infinite **n**
- $L^* = \{w_1 \dots w_n : w_i \in L, i = 1, 2, \dots, n, n \geq 1\}$
Justify: $n \geq 0$. **n**
- For any language L over an alphabet Σ , $L^+ = L \cup L^*$
Justify: $e \in L^*$ and $e \notin L^+$ **n**

12. For any language L over an alphabet Σ , $L^+ = L \cup L^*$.

Justify: Take L such that $e \notin L$. We get that $e \in L \cup L^*$ as $e \in L^*$ and $e \notin L^+$.

n

13. For any languages L_1, L_2, L over $\Sigma \neq \emptyset$

$$(L_1 \cup L_2) \circ L = L_1 \circ L \cup L_2 \circ L$$

Justify: proved in class

y

14. Regular language is a regular expression.

Justify: Regular Language is represented by the function \mathcal{L}

n

15. Regular expression $\alpha = (a \cup b)^* a$ defines a language

$$L = \{w \in \{a, b\}^* : w \text{ ends with } a\}$$

Justify: $\{a, b\}^* \{a\} = \Sigma^* \{a\} = \{w \in \{a, b\}^* : w \text{ ends with } a\}$.

y

PART 2: PROBLEMS

Chose ONE problem and solve it

Circle the problem you want us to correct

QUESTION 1

1. Let $A = \{(\{n, n+1\}, n) \in 2^N \times N : 1 \leq n \leq 3\}$. List all elements of A .
2. Let now $A = \{(\{n\}, n) \in 2^N \times N : 1 \leq n \leq n+1\}$. Prove that A is infinitely countable.

Solution

1. Let $A = \{(\{n, n+1\}, n) \in 2^N \times N : 1 \leq n \leq 3\}$. List all elements of A .

$$A = \{(\{n, n+1\}, n) \in 2^N \times N : n = 1, 2, 3\} = \{(\{1, 2\}, 1), (\{2, 3\}, 2), (\{3, 4\}, 3)\}$$

2. Let now $A = \{(\{n\}, n) \in 2^N \times N : 1 \leq n \leq n+1\}$. Prove that A is infinitely countable.

Solution

$$A = \{(\{n\}, n) \in 2^N \times N : 1 \leq n \leq n+1\} = \{(\{n\}, n) \in 2^N \times N : 1 \leq n\}$$

because $n \leq n+1$ for all $n \in N$.

The set $B = \{\{n\} : n \in N\}$ has the same cardinality as N by the function $f(n) = \{n\}$. $A = B \times N$ is a Cartesian product of two infinitely countable sets, and hence is also infinitely countable.

QUESTION 2 Let $\Sigma = \{a, b\}$. Let $L_1 \subseteq \Sigma^*$ be defined as follows:

$$L_1 = \{w \in \Sigma^* : \text{number of } b \text{ in } w \text{ is divisible three}\}$$

tion. Explain shortly your answer.

Solution: Write a regular expression α , such that $\mathcal{L}(\alpha) = L_1$. You can use shorthand nota

$$\alpha = a^* \cup (a^*ba^*ba^*ba^*)^*$$

Explanation: the part $a^*ba^*ba^*ba^*$ says that there must be 3 occurrences of b in L_1 . The part $(a^*ba^*ba^*ba^*)^*$ says that we the number of b 's is $3n$ for $n \geq 1$.

Observe that 0 is divisible by 3, so we need to add the case of 0 number of b 's ($n = 0$), i.e. words $e, a, aa, aaa, , \dots$. We do so by adding a^* to $(a^*ba^*ba^*ba^*)^*$

We can also do so by concatenating a^* with $(a^*ba^*ba^*ba^*)^*$, as $e \in (a^*ba^*ba^*ba^*)^*$ and get alternative solution

$$\alpha = a^*(a^*ba^*ba^*ba^*)^*$$