

CSE303 Q1 Solutions

PART 1: YES/NO QUESTIONS

Circle your answer and write a short justification (1pt) No justification- no points

1. $\{a, \{\emptyset\}, 2\} \cap 2^\emptyset = \emptyset$

Justify: $2^\emptyset = \{\emptyset\}$ and $\emptyset \notin \{a, \{\emptyset\}, 2\}$

y

2. The relation $R = \{(n, m) : n \in N \text{ and } m = 1\}$ is a function

Justify: R is a function defined by the formula is $R(n) = 1$ for all $n \in N$

y

3. If the A is uncountable, then $|A| = \mathcal{C}$

Justify: The set 2^R of all subsets of real numbers R is uncountable, but by Cantor Theorem we have that $|2^R| > |R| = \mathcal{C}$.

n

4. The set $A = \{x \in N : x \geq 0\}$ is finite

Justify: $A = \{x \in N : x \geq 0\} = N$ and N is countably infinite

n

5. The set $\{\emptyset, \{\emptyset\}, 1, 2\}$ is an alphabet

Justify: A is a finite set and any finite set is an alphabet by definition

y

6. Let $\Sigma = \{\emptyset\}$. There are uncountably many languages over Σ

Justify: $\Sigma \neq \emptyset$ as $\emptyset$ is element of Σ . Hence there are infinitely countably many elements in Σ^* and uncountably many subsets of Σ^* . In fact exactly as many as real numbers as $|2^{\Sigma^*}| = |R| = \mathcal{C}$.

y

7. For any languages L_1, L_2, L over $\Sigma \neq \emptyset$

$$(L_1 \cap L_2) \cup L = (L_1 \cup L) \cap (L_2 \cup L)$$

Justify: Languages are sets and this is the Law of Distributivity of union over intersection of sets

y

8. $L^* = \{w_1 \dots w_n : w_i \in L, i = 1, 2, \dots, n, n \geq 1\}$

Justify: This is definition of L^+ ; Kleene Star must have condition $n \geq 0$.

n

9. Regular language is a regular expression.

Justify: Regular language is *defined* by a regular expression.

n

10. For any language L over an alphabet Σ , $L^+ = L \cup L^*$.

Justify: holds only only when $e \in L$ as $e \in L^*$

n

PART 2

QUESTION 1

Given an alphabet $\Sigma = \{a, b\}$ and a regular expression $\alpha = a^*b \cup (a \cup b)^*$.

1. (3pts) Evaluate $L = \mathcal{L}(\alpha)$.

Solution

1. We evaluate

$$L = \mathcal{L}(a^*b \cup (a \cup b)^*) = \mathcal{L}(a^*)\mathcal{L}(b) \cup (\mathcal{L}(a) \cup \mathcal{L}(b))^* = \{a\}^*\{b\} \cup \{a, b\}^*$$

2. (4pts) Give a property describing the language L determined by α

Solution

Observe that $\{a\}^*\{b\} \subseteq \{a, b\}^*$ hence the language is

$$L = \{w : w \in \{a, b\}^*\} = \Sigma^*$$

QUESTION 2

Let $\Sigma = \{a, b\}$ and a language $L \subseteq \Sigma^*$ be defined as follows:

$$L = \{w \in \Sigma^* : w \text{ contains less than two } a\text{'s}\}$$

Write a regular expression α , such that $\mathcal{L}(\alpha) = L$. Use shorthand notation. **Explain** shortly your answer.

Solution

$$\alpha = b^* \cup b^*ab^*$$

Explanation

"w contains less than two a's" means "contains zero or one a"

b^* contains no of a (case n=0)

b^*ab^* contains one occurrence of a (case n=1)