

cse371/mat371  
LOGIC

Professor Anita Wasilewska

Fall 2018

# Chapter 7

## Introduction to Intuitionistic and Modal Logics

### Slides Set 2

### PART 4: Gentzen Sequent System **LI**

## Gentzen Sequent System **LI**

**G. Gentzen** formulated in **1935** a first **syntactically decidable** (in propositional case) **proof systems** for classical and intuitionistic logics

He proved their **equivalence** with their well established, respective **Hilbert style** formalizations

He **named** his **classical** system **LK** (**K** for Klassisch) and **intuitionistic** system **LI** (**I** for Intuitionistisch)

## Gentzen Sequent System **LI**

In order to prove the **completeness** of the system **LK** and to prove the **adequacy** of **LI** he **introduced** a special inference rule, called **cut** rule that **corresponds** to the **Modus Ponens** rule in **Hilbert** style proof systems

Then, as the **next step** he proved the now famous **Hauptsatz**, called in English the **Cut Elimination Theorem**

## Gentzen Sequent System LI

Gentzen original proof system LI is a particular case of his proof system LK for the classical logic

Both of them are presented in chapter 6 together with the original Gentzen's proof of the **Hauptsatz** for both, LK and LI proof systems

The elimination of the cut rule and the structure of other rules makes it possible to define effective automatic procedures for proof search, what is impossible in a case of the Hilbert style systems

## LI Sequents

The Gentzen system **LI** is defined as follows.

Let

$$SQ = \{ \Gamma \longrightarrow \Delta : \Gamma, \Delta \in \mathcal{F}^* \}$$

be the set of all **Gentzen sequents** built out of the formulas of the language

$$\mathcal{L} = \mathcal{L}_{\{ \cup, \cap, \Rightarrow, \neg \}}$$

and the additional **Gentzen** arrow symbol  $\longrightarrow$

We assume that all **LI** sequents are elements of a following subset **ISQ** of the set **SQ** of all sequents

$$ISQ = \{ \Gamma \longrightarrow \Delta : \Delta \text{ consists of at most one formula} \}$$

The set **ISQ** is called the set of all **intuitionistic sequents**; the **LI** sequents

## Axioms of LI

**Logical Axioms** of **LI** consist of any sequent from the set *ISQ* which contains a **formula** that appears on **both sides** of the sequent arrow  $\longrightarrow$ , i.e any sequent of the form

$$\Gamma, A, \Delta \longrightarrow A$$

for  $\Gamma, \Delta \in \mathcal{F}^*$

## Rules of Inference of LI

The set inference rules of LI is divided into **two groups** : the **structural rules** and the **logical rules**

There are three **Structural Rules** of LI: **Weakening**, **Contraction** and **Exchange**

**Weakening** structural rule

$$(weak \rightarrow) \frac{\Gamma \rightarrow \Delta}{A, \Gamma \rightarrow \Delta}$$

$$(\rightarrow weak) \frac{\Gamma \rightarrow}{\Gamma \rightarrow A}$$

**A** is called the **weakening formula**

**Remember** that  $\Delta$  contains **at most one formula**



## Rules of Inference of **LI**

**Contraction** structural rule

$$(contr \rightarrow) \frac{A, A, \Gamma \rightarrow \Delta}{A, \Gamma \rightarrow \Delta}$$

$A$  is called the **contraction formula**

**Remember** that  $\Delta$  contains **at most one formula**

The rule below is **not VALID** for **LI**; we list it as it is used in the classical case

$$(\rightarrow contr) \frac{\Gamma \rightarrow \Delta, A, A}{\Gamma \rightarrow \Delta, A}$$

## Rules of Inference of **LI**

**Exchange** structural rule

$$(exch \rightarrow) \frac{\Gamma_1, A, B, \Gamma_2 \rightarrow \Delta}{\Gamma_1, B, A, \Gamma_2 \rightarrow \Delta}$$

**Remember** that  $\Delta$  contains **at most one formula**

The rule below is **not VALID** for **LI**; we list it as it is used in the classical case

$$(\rightarrow exch) \frac{\Delta \rightarrow \Gamma_1, A, B, \Gamma_2}{\Delta \rightarrow \Gamma_1, B, A, \Gamma_2}.$$

## Rules of Inference of LI

### Logical Rules

#### Conjunction rules

$$(\wedge \rightarrow) \frac{A, B, \Gamma \rightarrow \Delta}{(A \wedge B), \Gamma \rightarrow \Delta},$$

$$(\rightarrow \wedge) \frac{\Gamma \rightarrow A ; \Gamma \rightarrow B}{\Gamma \rightarrow (A \wedge B)}$$

**Remember** that  $\Delta$  contains **at most one formula**

## Rules of Inference of LI

### Disjunction rules

$$(\rightarrow \cup)_1 \quad \frac{\Gamma \rightarrow A}{\Gamma \rightarrow (A \cup B)}$$

$$(\rightarrow \cup)_2 \quad \frac{\Gamma \rightarrow B}{\Gamma \rightarrow (A \cup B)}$$

$$(\cup \rightarrow) \quad \frac{A, \Gamma \rightarrow \Delta ; B, \Gamma \rightarrow \Delta}{(A \cup B), \Gamma \rightarrow \Delta}$$

**Remember** that  $\Delta$  contains **at most one formula**

## Rules of Inference of LI

### Implication rules

$$(\rightarrow \Rightarrow) \frac{A, \Gamma \rightarrow B}{\Gamma \rightarrow (A \Rightarrow B)}$$

$$(\Rightarrow \rightarrow) \frac{\Gamma \rightarrow A ; B, \Gamma \rightarrow \Delta}{(A \Rightarrow B), \Gamma \rightarrow \Delta}$$

**Remember** that  $\Delta$  contains **at most one formula**

## Gentzen System LI

### Negation rules

$$(\neg \rightarrow) \frac{\Gamma \rightarrow A}{\neg A, \Gamma \rightarrow}$$

$$(\rightarrow \neg) \frac{A, \Gamma \rightarrow}{\Gamma \rightarrow \neg A}$$

We define the Gentzen system LI as

$$\mathbf{LI} = (\mathcal{L}, ISQ, LA, \text{Structural rules}, \text{Logical rules})$$

## LI Completeness

The completeness of the **cut-free LI** follows directly from **LI Hauptzatz** proved in chapter 6 and the **intuitionistic completeness** (Mostowski 1948)

### Completeness of LI

For any sequent  $\Gamma \rightarrow \Delta \in ISQ$ ,

$$\vdash_{LI} \Gamma \rightarrow \Delta \quad \text{if and only of} \quad \models_I \Gamma \rightarrow \Delta$$

In particular, for any formula  $A$ ,

$$\vdash_{LI} A \quad \text{if and only of} \quad \models_I A$$

## Intuitionistic Disjunction

The particular form the following theorem was stated without the proof by Gödel in 1931

The theorem proved by Gentzen in 1935 via **Hauptsatz** and we follow his proof

### Intuitionistically Derivable Disjunction

For any formulas  $A, B \in \mathcal{F}$ ,

$$\vdash_{LI} (A \cup B) \quad \text{if and only if} \quad \vdash_{LI} A \quad \text{or} \quad \vdash_{LI} B$$

In particular, a disjunction  $(A \cup B)$  is intuitionistically **provable** in any proof system  $I$  if and only if either  $A$  or  $B$  is intuitionistically **provable** in  $I$



## Intuitionistic Disjunction

### Proof of

$\vdash_{LI} (A \cup B)$  if and only if  $\vdash_{LI} A$  or  $\vdash_{LI} B$

Assume  $\vdash_{LI} (A \cup B)$

This equivalent to  $\vdash_{LI} \rightarrow (A \cup B)$

The **last** step in the proof of  $\rightarrow (A \cup B)$  in **LI** must be the application of the rule  $(\rightarrow \cup)_1$  to the sequent  $\rightarrow A$ , or the application of the rule  $(\rightarrow \cup)_2$  to the sequent  $\rightarrow B$

There is no other possibilities

We have proved that  $\vdash_{LI} (A \cup B)$  implies  $\vdash_{LI} A$  or  $\vdash_{LI} B$

The **inverse** implication is obvious by respective applications of rules  $(\rightarrow \cup)_1$  or  $(\rightarrow \cup)_2$  to the sequents  $\rightarrow A$  or  $\rightarrow B$

## Decomposition Trees in LI

## Decomposition Trees in LI

**Search for proofs** in **LI** is a much more complicated process than the one in classical logic systems **RS** or **GL** defined in chapter 6

Here, as in any other **Gentzen style** proof system, proof search **procedure** consists of building the **decomposition** trees

### Remark 1

In **RS** the **decomposition** tree  $T_A$  of any formula  $A$  is always **unique**

### Remark 2

In **GL** the "blind search" defines, for any formula  $A$  a **finite** number of **decomposition** trees,

Nevertheless, it can be proved that the search can be reduced to examining only **one** of them, due to the **absence** of structural rules

## Decomposition Trees in LI

### Remark 3

In LI the **structural rules** play a **vital role** in the proof construction and hence, in the proof search

The fact that a given **decomposition tree** ends with an **non-axiom leaf** **does not** always imply that the proof **does not** exist

It might only imply that our **search strategy** was **not good**

The problem of **deciding** whether a given formula **A** **does**, or **does not** have a proof in LI becomes more **complex** than in the case of Gentzen system for **classical** logic

## Decomposition Trees in LI

Before we define a **heuristic method** of **searching** for proof and **deciding** whether such a proof **exists** or **not** we make some observations

### Observation 1

**Logical rules** of **LI** are similar to those in Gentzen type **classical** formalizations we already examined in previous chapters in a sense that each of them **introduces** a logical **connective**

## Decomposition Trees in LI

### Observation 2

The process of searching for a proof is a **decomposition** process in which we use the **inverse** of logical and structural rules as **decomposition** rules

For **example** the implication rule:

$$(\rightarrow\Rightarrow) \frac{A, \Gamma \rightarrow B}{\Gamma \rightarrow (A \Rightarrow B)}$$

becomes an implication **decomposition** rule (we use the same name  $(\rightarrow\Rightarrow)$  in both cases)

$$(\rightarrow\Rightarrow) \frac{\Gamma \rightarrow (A \Rightarrow B)}{A, \Gamma \rightarrow B}$$

## Decomposition Trees in LI

### Observation 3

We write proofs as **trees**, so the **proof search** process is a process of building **decomposition** trees

To **facilitate** the process we write the **decomposition** rules in a **tree** decomposition form as follows

$$\Gamma \longrightarrow (A \Rightarrow B)$$

$$| (\rightarrow \Rightarrow)$$

$$A, \Gamma \longrightarrow B$$

## Decomposition Trees in LI

The two premisses rule  $(\Rightarrow \rightarrow)$  written as the tree decomposition rule becomes

$$\frac{(A \Rightarrow B), \Gamma \rightarrow}{\bigwedge (\Rightarrow \rightarrow)} \frac{\Gamma \rightarrow A \quad B, \Gamma \rightarrow}{}$$



## Decomposition Trees in LI

The structural **weakening** rule written as the **decomposition** rule is

$$(\rightarrow \text{weak}) \frac{\Gamma \rightarrow A}{\Gamma \rightarrow}$$

We write it in a **tree decomposition** form as

$$\Gamma \rightarrow A$$

$$| (\rightarrow \text{weak})$$

$$\Gamma \rightarrow$$

## Decomposition Trees in LI

We define the notion of **decomposable** and **indecomposable** formulas and sequents as follows

**Decomposable formula** is any formula of the **degree  $\geq 1$**

**Decomposable sequent** is any sequent that contains a **decomposable** formula

**Indecomposable formula** is any formula of the **degree 0**  
i.e. is any **propositional variable**

## Decomposition Trees in LI

### Remark

In a case of **formulas** written with use of capital letters **A, B, C, .. etc** , we treat these letters as propositional **variables** , i.e. as **indecomposable formulas**

**Indecomposable sequent** is a sequent formed from **indecomposable formulas** only.

## Decomposition Trees in LI

### Decomposition Tree Construction (1)

Given a formula  $A$  we construct its **decomposition** tree  $T_A$  as follows

**Root** of the tree  $T_A$  is the sequent  $\longrightarrow A$

Given a **node**  $n$  of the tree we identify a **decomposition** rule **applicable** at this node and write its **premisses** as the **leaves** of the **node**  $n$

We **stop** the decomposition **process** when we obtain an **axiom** or **all leaves** of the tree are **indecomposable**

## Decomposition Trees in LI

### Observation 4

The decomposition tree  $T_A$  obtained by the **Construction (1)** most often **is not unique**

### Observation 5

The fact that we **find** a decomposition tree  $T_A$  with a **non-axiom** leaf **does not** mean that  $\not\vdash_{LI} A$

This is due to the **role** of **structural rules** in **LI** and will be discussed later

## Proof Search Examples

## Examples

We perform **proof search** and **decide** the existence of proofs in **LI** for a given formula  $A \in \mathcal{F}$  by constructing its **decomposition trees**  $T_A$

We examine here some **examples** to show the **complexity** of the problem

### Reminder

In the following and **similar** examples when building the decomposition trees for formulas representing **general schemas** we treat the capital letters  $A, B, C, D...$  as **propositional** variables, i.e. as **indecomposable** formulas

## Examples

### Example 1

Determine] whether

$$\vdash_{\mathbf{LI}} ((\neg A \cap \neg B) \Rightarrow \neg(A \cup B))$$

**Observe** that

If we find a decomposition tree of  $A$  in  $\mathbf{LI}$  such that **all its leaves are axiom**, we have a proof, i.e.

$$\vdash_{\mathbf{LI}} A$$

If **all possible** decomposition trees have a **non-axiom leaf** then the proof of  $A$  in  $\mathbf{LI}$  does not exist, i.e.

$$\not\vdash_{\mathbf{LI}} A$$



## Examples

Consider the following decomposition tree  $T1_A$

$$\rightarrow ((\neg A \cap \neg B) \Rightarrow \neg(A \cup B))$$

$$| (\rightarrow \Rightarrow)$$

$$(\neg A \cap \neg B) \rightarrow \neg(A \cup B)$$

$$| (\rightarrow \neg)$$

$$(\neg A \cap \neg B), (A \cup B) \rightarrow$$

$$| (\cap \rightarrow)$$

$$\neg A, \neg B, (A \cup B) \rightarrow$$

$$| (\neg \rightarrow)$$

$$\neg B, (A \cup B) \rightarrow A$$

$$| (\rightarrow \text{weak})$$

$$\neg B, (A \cup B) \rightarrow$$

$$| (\neg \rightarrow)$$

$$(A \cup B) \rightarrow B$$

$$\bigwedge (U \rightarrow)$$

$$A \rightarrow B$$

*non - axiom*

$$B \rightarrow B$$

*axiom*

## Examples

The tree  $T1_A$  has a **non-axiom** leaf, so it **does not** constitute a proof in **LI**

Observe that the **decomposition** tree in **LI** is not always **unique**

Hence the existence of a **non-axiom** leaf **does not** yet prove that the **proof** of **A** does not **exist**

Consider the following decomposition tree  $T2_A$

$$\rightarrow ((\neg A \cap \neg B) \Rightarrow (\neg(A \cup B)))$$

$$| (\rightarrow \Rightarrow)$$

$$(\neg A \cap \neg B) \rightarrow \neg(A \cup B)$$

$$| (\rightarrow \neg)$$

$$(A \cup B), (\neg A \cap \neg B) \rightarrow$$

$$| (exch \rightarrow)$$

$$(\neg A \cap \neg B), (A \cup B) \rightarrow$$

$$| (\cap \rightarrow)$$

$$\neg A, \neg B, (A \cup B) \rightarrow$$

$$| (exch \rightarrow)$$

$$\neg A, (A \cup B), \neg B \rightarrow$$

$$| (exch \rightarrow)$$

$$(A \cup B), \neg A, \neg B \rightarrow$$

$$\bigwedge (\cup \rightarrow)$$

$$A, \neg A, \neg B \rightarrow$$

$$| (exch \rightarrow)$$

$$\neg A, A, \neg B \rightarrow$$

$$| (\neg \rightarrow)$$

$$A, \neg B \rightarrow A$$

*axiom*

$$B, \neg A, \neg B \rightarrow$$

$$| (exch \rightarrow)$$

$$B, \neg B, \neg A \rightarrow$$

$$| (exch \rightarrow)$$

$$\neg B, B, \neg A \rightarrow$$

$$| (\neg \rightarrow)$$

$B, \neg A \rightarrow B$ ; *axiom*

## Examples

All leaves of  $T_{2A}$  are axioms

This means that the tree  $T_{2A}$  is a **proof** of  $A$  in  $LI$

We hence proved that

$$\vdash_{LI} ((\neg A \cap \neg B) \Rightarrow \neg(A \cup B))$$

## Examples

**Example 2:** Show that

1.  $\vdash_{LI} (A \Rightarrow \neg\neg A)$

2.  $\not\vdash_{LI} (\neg\neg A \Rightarrow A)$

**Solution of 1.**

We construct **some**, or **all decomposition** trees of

$$\longrightarrow (A \Rightarrow \neg\neg A)$$

A tree  $T_A$  that **ends** with **all** leaves being **axioms** is a proof of  $A$  in **LI**

## Examples

We construct  $T_A$  as follows

$$\longrightarrow (A \Rightarrow \neg\neg A)$$

$$| (\longrightarrow \Rightarrow)$$

$$A \longrightarrow \neg\neg A$$

$$| (\longrightarrow \neg)$$

$$\neg A, A \longrightarrow$$

$$| (\neg \longrightarrow)$$

$$A \longrightarrow A$$

*axiom*

All leaves of  $T_A$  are **axioms** so we found the **proof**

We **do not** need to construct any other decomposition trees.

## Examples

### Solution of 2.

In order to prove that

$$\not\vdash_{LI} (\neg\neg A \Rightarrow A)$$

we have to construct **all decomposition** trees of

$$\longrightarrow (\neg\neg A \Rightarrow A)$$

and show that **each** of them has a **non-axiom** leaf

## Examples

Here is **T1<sub>A</sub>**

$$\longrightarrow (\neg\neg A \Rightarrow A)$$

$$| (\longrightarrow \Rightarrow)$$

*one of 2 choices*

$$\neg\neg A \longrightarrow A$$

$$| (\longrightarrow \text{weak})$$

*one of 3 choices*

$$\neg\neg A \longrightarrow$$

$$| (\neg \longrightarrow)$$

*one of 3 choices*

$$\longrightarrow \neg A$$

$$| (\longrightarrow \neg)$$

*one of 2 choices*

$$A \longrightarrow$$

*non - axiom*



# Here is **T2<sub>A</sub>**

$$\rightarrow (\neg\neg A \Rightarrow A)$$

| ( $\rightarrow\Rightarrow$ ) *one of 2 choices*

$$\neg\neg A \rightarrow A$$

| (*contr*  $\rightarrow$ ) *second of 2 choices*

$$\neg\neg A, \neg\neg A \rightarrow A$$

| ( $\rightarrow$  *weak*) *first of 2 choices*

$$\neg\neg A, \neg\neg A \rightarrow$$

| ( $\neg\rightarrow$ ) *first of 2 choices*

$$\neg\neg A \rightarrow \neg A$$

| ( $\rightarrow\neg$ ) *one of 2 choices*

$$A, \neg\neg A \rightarrow$$

| (*exch*  $\rightarrow$ ) *one of 2 choices*

$$\neg\neg A, A \rightarrow$$

| ( $\neg\rightarrow$ ) *one of 2 choices*

$$A \rightarrow \neg A$$

| ( $\rightarrow\neg$ ) *first of 2 choices*

$$A, A \rightarrow$$

*non - axiom*

## Structural Rules

We can see from the above **decomposition** trees that the "blind" construction of all possible trees only leads to more complicated trees

This is due to the presence of structural rules

The "blind" application of the rule (*contr*  $\rightarrow$ ) gives always an infinite number of **decomposition** trees

In order to decide that none of them will produce a proof we need some **extra knowledge** about patterns of their construction, or just simply about the number of useful of application of **structural rules**

## Structural Rules

In this case we can just make an "external" **observation** that the our first tree  $T_{1A}$  is in a sense a **minimal one**

It means that all **other trees** would only **complicate** this one in an **inessential way**, i.e. the we will **never produce** a tree with all **axioms leaves**

One can formulate a **deterministic procedure** giving a finite number of trees, but the proof of its **correctness** is needed and that requires some **extra knowledge**

Within the scope of this book we accept the **"external explanation** as a **sufficient solution**

## Structural Rules

As we can see from the above examples the **structural rules** and especially the (*contr*  $\rightarrow$ ) rule **complicates** the proof searching task.

Both **Gentzen type** proof systems **RS** and **GL** from the previous chapter **don't contain** the structural rules

They also are as we have proved, **complete** with respect to classical semantics.

The **original Gentzen** system **LK** which does contain the structural rules is also, as proved by Gentzen, **complete**

## Structural Rules

Hence **all three** classical proof system **RS, GL, LK** are **equivalent**

This proves that the **structural rules** can be **eliminated** from the system **LK**

A natural question of **elimination** of **structural rules** from the system **LI** arises

The following **example** illustrates the **negative answer**

## Examples

### Example 3

We know that for any formula  $A \in \mathcal{F}$ ,

$$\models A \quad \text{if and only if} \quad \vdash_I \neg\neg A$$

where  $\models A$  means that  $A$  is **classical** tautology

$\vdash_I A$  means that  $A$  is **Intuitionistically provable** in any intuitionistically **complete** proof system  $I$

The system **LI** is intuitionistically **complete** so have that for any formula  $A \in \mathcal{F}$ ,

$$\models A \quad \text{if and only if} \quad \vdash_{LI} \neg\neg A$$

## Examples

Obviously  $\models (\neg\neg A \Rightarrow A)$ , so we must have that

$$\vdash_{LI} \neg\neg(\neg\neg A \Rightarrow A)$$

We are going to prove now that the rule  $(\text{contr} \rightarrow)$  is **essential** to the **existence** of the proof  $\neg\neg(\neg\neg A \Rightarrow A)$

It means that  $\neg\neg(\neg\neg A \Rightarrow A)$  **is not provable** without the rule  $(\text{contr} \rightarrow)$

The following decomposition tree  $\mathbf{T}_A$  is a proof of  $\neg\neg(\neg\neg A \Rightarrow A)$  **with use** of the rule  $(\text{contr} \rightarrow)$

# Examples

$$\rightarrow \neg\neg(\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \neg)$$

$$\neg(\neg\neg A \Rightarrow A) \rightarrow$$

$$| (\text{contr} \rightarrow)$$

$$\neg(\neg\neg A \Rightarrow A), \neg(\neg\neg A \Rightarrow A) \rightarrow$$

$$| (\neg \rightarrow)$$

$$\neg(\neg\neg A \Rightarrow A) \rightarrow (\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \Rightarrow)$$

$$\neg(\neg\neg A \Rightarrow A), \neg\neg A \rightarrow A$$

$$| (\rightarrow \text{weak})$$

$$\neg(\neg\neg A \Rightarrow A), \neg\neg A \rightarrow$$

$$| (\text{exch} \rightarrow)$$

$$\neg\neg A, \neg(\neg\neg A \Rightarrow A) \rightarrow$$

$$| (\neg \rightarrow)$$

$$\neg(\neg\neg A \Rightarrow A) \rightarrow \neg A$$

$$| (\rightarrow \neg)$$

$$A, \neg(\neg\neg A \Rightarrow A) \rightarrow$$

$$| (\text{exch} \rightarrow)$$

$$\neg(\neg\neg A \Rightarrow A), A \rightarrow$$

$$| (\neg \rightarrow)$$

$$A \rightarrow (\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \Rightarrow)$$



## Contraction Rule

Assume now that the rule (*contr*  $\rightarrow$ ) is **not** available. All **possible** decomposition trees are as follows

Tree **T1<sub>A</sub>**

$$\rightarrow \neg\neg(\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \neg)$$

$$\neg(\neg\neg A \Rightarrow A) \rightarrow$$

$$| (\neg \rightarrow)$$

$$\rightarrow (\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \Rightarrow)$$

$$\neg\neg A \rightarrow A$$

$$| (\rightarrow \textit{weak})$$

$$\neg\neg A \rightarrow$$

$$| (\neg \rightarrow)$$

$$\rightarrow \neg A$$

$$| (\rightarrow \neg)$$

$$A \rightarrow$$

*non - axiom*

## Contraction Rule

The next is **T2<sub>A</sub>**

$$\longrightarrow \neg\neg(\neg\neg A \Rightarrow A)$$

$$| (\longrightarrow \neg)$$

$$\neg(\neg\neg A \Rightarrow A) \longrightarrow$$

$$| (\neg \longrightarrow)$$

$$\longrightarrow (\neg\neg A \Rightarrow A)$$

$$| (\longrightarrow \textit{weak})$$

$\longrightarrow$

*non - axiom*

## Contraction Rule

The next is **T3<sub>A</sub>**

$$\longrightarrow \neg\neg(\neg\neg A \Rightarrow A)$$

| ( $\longrightarrow$  weak)

$\longrightarrow$

*non - axiom*

## Contraction Rule

The last one is **T4<sub>A</sub>**

$$\rightarrow \neg\neg(\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \neg)$$

$$\neg(\neg\neg A \Rightarrow A) \rightarrow$$

$$| (\neg \rightarrow)$$

$$\rightarrow (\neg\neg A \Rightarrow A)$$

$$| (\rightarrow \Rightarrow)$$

]

$$\neg\neg A \rightarrow A$$

$$| (\rightarrow \text{weak})$$

$$\neg\neg A \rightarrow$$

$$| (\neg \rightarrow)$$

$$\rightarrow \neg A$$

$$| (\rightarrow \text{weak})$$

$\rightarrow$

*non - axiom*

## Contraction Rule

We have considered all **possible** decomposition trees that **do not** involve the contraction rule (*contr*  $\longrightarrow$ ) and **none** of them was a proof

This shows that the formula

$$\neg\neg(\neg\neg A \Rightarrow A)$$

**is not provable** in **LI** without (*contr*  $\longrightarrow$ ) rule, i.e. that we proved the following

### Fact

The contraction rule (*contr*  $\longrightarrow$ ) **can not** be **eliminated** from **LI**

## Proof Search Heuristic Method

## Proof Search Heuristic Method

Before we define a heuristic method of searching for proof in **LI** let's make some **additional** observations to the already made **observations 1-5**

### Observation 6

The **goal** of constructing the decomposition tree is to **obtain axioms** or **indecomposable** leaves

With respect to this goal the **use logical** decomposition rules has **a priority** over the use of the **structural** rules

We use this information while describing the proof search **heuristic**

## Proof Search Heuristic Method

### Observation 7

All logical decomposition rules ( $\circ \rightarrow$ ), where  $\circ$  denotes any connective, must have a formula we want to decompose as the **first formula** at the decomposition node

It means that if we want to **decompose** a formula  $\circ A$  the node must have a form  $\circ A, \Gamma \rightarrow \Delta$

**Remember:** order of decomposition is important

Also sometimes **it is necessary** to decompose a **formula within the sequence  $\Gamma$  first**, before decomposing  $\circ A$  in order to **find** a proof



## Proof Search Heuristic Method

For example, consider two nodes

$$n_1 = \neg\neg A, (A \cap B) \longrightarrow B$$

and

$$n_2 = (A \cap B), \neg\neg A \longrightarrow B$$

We are going to see that the results of decomposing  $n_1$  and  $n_2$  **differ dramatically**

Let's decompose the node  $n_1$

Observe that the only way to be able to decompose the formula  $\neg\neg A$  is to use the rule ( $\rightarrow$  *weak*) as a **first step**

The **two possible** decomposition trees that **starts at the node**  $n_1$  are as follows

## Proof Search Heuristic Method

### First Tree

**T1**<sub>m1</sub>

$\neg\neg A, (A \cap B) \longrightarrow B$

| ( $\rightarrow$  weak)

$\neg\neg A, (A \cap B) \longrightarrow$

| ( $\neg \rightarrow$ )

$(A \cap B) \longrightarrow \neg A$

| ( $\cap \rightarrow$ )

$A, B \longrightarrow \neg A$

| ( $\rightarrow \neg$ )

$A, A, B \longrightarrow$

*non - axiom*

## Proof Search Heuristic Method

### Second Tree

**T2<sub>m1</sub>**

$$\neg\neg A, (A \cap B) \longrightarrow B$$

| ( $\rightarrow$  weak)

$$\neg\neg A, (A \cap B) \longrightarrow$$

| ( $\neg \rightarrow$ )

$$(A \cap B) \longrightarrow \neg A$$

| ( $\rightarrow \neg$ )

$$A, (A \cap B) \longrightarrow$$

| ( $\cap \rightarrow$ )

$$A, A, B \longrightarrow$$

*non - axiom*

## Proof Search Heuristic Method

Let's now decompose the node  $n_2$

Observe that following our **Observation 6** we **start** by decomposing the formula  $(A \cap B)$  by the use of the rule  $(\cap \rightarrow)$  as the **first step**

A decomposition tree that starts at the node  $n_2$  is as follows

$T_{n_2}$

$$(A \cap B), \neg\neg A \longrightarrow B$$

$$| (\cap \rightarrow)$$

$$A, B, \neg\neg A \longrightarrow B$$

*axiom*

This proves that the node  $n_2$  is **provable** in **LI**, i.e.

$$\vdash_{LI} (A \cap B), \neg\neg A \longrightarrow B$$

## Proof Search Heuristic Method

### Observation 8

The use of **structural rules** is **important** and **necessary** while we search for proofs

Nevertheless we have to **use them** on the **"must" basis** and set up some **guidelines** and **priorities** for their use

For example, the use of **weakening rule** **discharges** the **weakening formula**, and hence we might **lose an information** that may be **essential** to finding the **proof**

We should use the **weakening rule** only when it is **absolutely necessary** for the next decomposition steps

## Proof Search Heuristic Method

Hence, the use of weakening rule ( $\rightarrow$  *weak*) **can**, and **should be restricted** to the cases when it leads to **possibility** of the future use of the **negation rule** ( $\neg \rightarrow$ )

This was the case of the decomposition tree **T1**<sub>n<sub>1</sub></sub>

We used the rule ( $\rightarrow$  *weak*) as an **necessary step**, but it **discharged** too much information and we **didn't get a proof**, when **proof on this node existed**

## Proof Search Heuristic Method

Here is such a proof

**T3**<sub>n<sub>1</sub></sub>

$$\neg\neg A, (A \cap B) \longrightarrow B$$

| (*exch*  $\longrightarrow$ )

$$(A \cap B), \neg\neg A \longrightarrow B$$

| ( $\cap \longrightarrow$ )

$$A, B, \neg\neg A \longrightarrow B$$

*axiom*

## Proof Search Heuristic Method

### Method

For any  $A \in \mathcal{F}$  we construct the set of decomposition trees  $\mathbf{T}_{\rightarrow A}$  following the rules below.

1. Use first **logical rules** where applicable.
2. Use (*exch*  $\rightarrow$ ) rule to decompose, via **logical rules**, as many formulas on the left side of  $\rightarrow$  as possible

**Remember** that the **order of decomposition** matters! so you have to cover different choices

3. Use ( $\rightarrow$  *weak*) only on a "**must**" basis and in connection with ( $\neg \rightarrow$ ) rule
4. Use (*contr*  $\rightarrow$ ) rule as the **last recourse** and only to formulas that contain  $\neg$  or  $\Rightarrow$  as a main connective
5. Let's call a formula  $A$  to which we apply (*contr*  $\rightarrow$ ) rule a **a contraction formula**
6. The only contraction formulas are formulas containing  $\neg$  between their logical connectives



## Proof Search Heuristic Method

7. Within the process of construction of all possible trees use (*contr*  $\rightarrow$ ) rule **only** to **contraction formulas**
8. Let  $C$  be a **contraction formula** appearing on a node  $n$  of the decomposition tree of  $T \rightarrow A$

For any **contraction formula**  $C$ , any node  $n$ , we apply (*contr*  $\rightarrow$ ) rule the the formula  $C$  **at most** as many times as the number of sub-formulas of  $C$

If we **find** a tree with **all axiom leaves** we have a **proof**, i.e.

$$\vdash_{LI} A$$

If **all trees** (finite number) have a **non-axiom leaf** we have proved that proof of  $A$  **does not exist**, i.e.

$$\not\vdash_{LI} A$$