

cse541
LOGIC for Computer Science

Professor Anita Wasilewska

LECTURE 4

Chapter 4

GENERAL PROOF SYSTEMS

PART 1: Introduction- Intuitive definitions

PART 2: Formal Definition of a Proof System

PART 3: Formal Proofs and Simple Examples

PART 4: Consequence, Soundness and Completeness

PART 5: Decidable and Syntactically Decidable Proof Systems

PART 1: General Introduction

Proof Systems - Intuitive Definition

Proof systems are built to prove, it means to **construct formal proofs** of statements formulated in a given **language**

First component of any **proof system** is hence its **formal language** \mathcal{L}

Proof systems are **inference machines** with statements called **provable statements** being their **final products**

Semantical Link

The **starting points** of the **inference machine** of a proof system **S** are called its **axioms**

We distinguish two kinds of axioms: **logical axioms** **LA** and **specific axioms** **SA**

Semantical link: we usually build a **proof systems** for a given **language** and its **semantics** i.e. for a **logic defined semantically**

Semantical Link

We always choose as a set of **logical axioms** **LA** some **subset of tautologies**, under a given **semantics**

We will **consider here** only proof systems with **finite sets** of **logical** or **specific axioms**, i.e we will examine only **finitely axiomatizable** proof systems

Semantical Link

We can, and we often do, consider **proof systems** with languages **without yet established semantics**

In this case the **logical axioms LA** serve as description of **tautologies** under a **future semantics** yet to be built

Logical axioms LA of a proof system **S** are hence not only **tautologies** under an established **semantics**, but they can also guide us how to **define a semantics** when it is yet **unknown**

Specific Axioms

The **specific axioms SA** consist of statements that describe a specific knowledge of an universe we want to use the proof system **S** to prove facts about

Specific axioms SA are not universally true

Specific axioms SA are **true only** in the universe we are interested to **describe** and **investigate** by the use of the proof system **S**

Formal Theory

Given a **proof system** S with **logical axioms** LA

Specific axioms SA of the proof system S is any finite set of formulas that **are not tautologies**, and hence they are always disjoint with the set of **logical axioms** LA of S

The **proof system** S with added set of **specific axioms** SA is called a **formal theory** based on S

Inference Machine

The **inference machine** of a proof system **S** is defined by a **finite set** of **inference rules**

The **inference rules** describe the way we are allowed to **transform** the information within the system with **axioms** as a starting point

We depict it **informally** on the next slide

Inference Machine

AXIOMS



RULES applied to AXIOMS



RULES applied to any expressions above



Provable formulas

Semantical Link

Semantical link:

Rules of inference of a system **S** have to **preserve the truthfulness** of what they are being used to prove

The notion of **truthfulness** is always defined by a given **semantics M**

Rules of inference that **preserve the truthfulness** are called **sound rules** under a given **semantics M**

Rules of inference can be **sound** under one semantics and **not sound** under another

Soundness Theorem

Goal 1

When developing a proof system **S** the first goal is prove the following theorem about it and its semantics **M**

Soundness Theorem

For any formula **A** of the language of the system **S**

If a formula **A** is **provable** from **logical axioms** **LA** of **S** only, then **A** is a **tautology** under the **semantics** **M**

Propositional Proof Systems

We discuss here first only proof systems for **propositional languages** and call them **proof systems** for different **propositional logics**

Remember

The notion of **soundness** is connected with a given **semantics**

A proof system **S** can be **sound** under **one semantics**, and **not sound** under the **other**

For example a set of axioms and rules **sound under classical logic semantics** might **not be sound** under **\perp logic semantics**, or **K logic semantics**, or others

Completeness of the Proof Systems

In general there are **many** proof systems that are **sound** under a given **semantics**, i.e. there are many **sound** proof systems for a given **logic** semantically defined

Given a proof system **S** with **logical axioms LA** that is **sound** under a **semantics M**.

Notation

Denote by T_M the set of all **tautologies** defined by the semantics **M**, i.e. we have that

$$T_M = \{A \in \mathcal{F} : \models_M A\}$$

Completeness Property

A **natural question** arises:

Are all **tautologies** i.e formulas $A \in T_M$ **provable** in the system **S** ??

We assume that we have already proved that **S** is **sound** under the semantics **M**

The **positive answer** to this question is called **completeness property** of the system **S** .

Completeness Theorem

Goal 2

Given for a **sound** proof system **S** under its semantics **M**, our the second goal is to prove the following theorem about **S**

Completeness Theorem

For any formula **A** of the language of **S**

A is provable in **S** iff **A** is a **tautology** under the semantics **M**

We write the **Completeness Theorem** *symbolically* as

$$\vdash_S A \text{ iff } \models_M A$$

Completeness Theorem is composed of two parts:

Soundness Theorem and the **Completeness Part** that proves the **completeness property** of a sound proof system

Proving Soundness and Completeness

Proving the **Soundness Theorem** for **S** under a semantics **M** is usually a straightforward and not a very difficult task

We **first prove** that all **logical axioms LA** are **tautologies**, and then we **prove** that all **inference rules** of the system **S** **preserve** the notion of the truth

Proving the **completeness part** of the **Completeness Theorem** is always a crucial, difficult and sometimes impossible task

OUR PLAN

We will study **two proofs** of the **Completeness Theorem** for **classical propositional** proof system in **Chapter 5**

We will present a **constructive** proofs of **Completeness Theorem** for two different **Gentzen style** automated theorem proving systems for **classical Logic** in **Chapter 6**

We discuss the **Intuitionistic Logic** in **Chapter 7**

Predicate Logics are discussed **Chapters 8, 9, 10, 11**

PART 2
PROOF SYSTEMS: Formal Definitions

Proof System S

In this section we present **formal definitions** of the following notions

Proof system S

Formal proof from **logical axioms** in a proof system S

Formal proof from **specific axioms** in a proof system S

Formal Theory based on a proof system S

We also give **examples** of different simple **proof systems**

Components: Language

Language \mathcal{L} of a **proof system** \mathbf{S} is any formal language \mathcal{L}

$$\mathcal{L} = (\mathcal{A}, \mathcal{F})$$

We assume as before that both sets \mathcal{A} and \mathcal{F} are enumerable, i.e. we deal here with **enumerable languages**

The **Language** \mathcal{L} can be **propositional** or **first order** (predicate) but we discuss **propositional languages** first

Components: Expressions

Expressions \mathcal{E} of a proof system \mathbf{S}

Given a set \mathcal{F} of well formed formulas of the language \mathcal{L} of the system \mathbf{S}

We often **extend** the set \mathcal{F} to some set \mathcal{E} of **expressions** build out of the language \mathcal{L} and some **extra symbols**, if needed

In this case all other **components** of \mathbf{S} are also defined on basis of elements of the set of **expressions** \mathcal{E}

In particular, and **most common case** we have that $\mathcal{E} = \mathcal{F}$

Expressions Examples

Automated theorem proving systems usually use as their **basic components** different sets of **expressions** build out of **formulas** of the language \mathcal{L}

In **Chapters 6 and 10** we consider **finite sequences of formulas** instead of formulas, as **basic expressions** of the proof systems **RS** and **RQ**

We also present there proof systems that use yet other kind of **expressions**, called original **Gentzen sequents** or their modifications

Some systems use yet **other expressions** such as **clauses**, **sets of clauses**, or **sets of formulas**, others use **yet still different** expressions

Semantical Link

We always have to **extend** a given semantics **M** for the **language** \mathcal{L} of the system **S** to the set \mathcal{E} of all **expression** of the system **S**

Sometimes, like in case of **Resolution** based **proof systems** we have also to **prove** a **semantic equivalency** of new created expressions \mathcal{E} (sets of clauses in Resolution case) with appropriate formulas of \mathcal{L}

Example

For example, in the automated theorem proving system **RS** presented in Chapter 6 the basic expressions \mathcal{E} are **finite sequences** of formulas of $\mathcal{L} = \mathcal{L}_{\{\neg, \wedge, \vee, \Rightarrow\}}$.

We **extend** our classical semantics for \mathcal{L} to the set \mathcal{F}^* of all **finite sequences** of formulas as follows:

For any $v : VAR \rightarrow \{F, T\}$ and any $\Delta \in \mathcal{F}^*$, $\Delta = A_1, A_2, \dots, A_n$, we put

$$\begin{aligned} v^*(\Delta) &= v^*(A_1, A_2, \dots, A_n) \\ &= v^*(A_1) \cup v^*(A_2) \cup \dots \cup v^*(A_n) \end{aligned}$$

i.e. in a shorthand notation

$$\Delta \equiv (A_1 \cup A_2 \cup \dots \cup A_n)$$

Components: Logical Axioms

Logical axioms LA of S form a **non-empty** subset of the set \mathcal{E} of **expressions** of the proof system S , i.e.

$$LA \subseteq \mathcal{E}$$

In particular, LA is a non-empty subset of **formulas**, i.e.

$$LA \subseteq \mathcal{F}$$

We **assume here** that the set LA of **logical axioms** is always **finite**, i.e. that we consider here **finitely axiomatizable** systems

In general, **we assume** that the set LA is **primitively recursive** i.e. that there is an effective procedure to determine whether a given expression $E \in \mathcal{E}$ **is** or **is not** in LA

Components: Axioms

Semantical link

Given a semantics \mathbf{M} for \mathcal{L} and its **extension** to the set \mathcal{E} of all expressions

We extend the notion of **tautology** to the expressions and write

$$\models_{\mathbf{M}} E$$

to denote that the **expression** $E \in \mathcal{E}$ is a **tautology** under semantics \mathbf{M} and we put

$$\mathbf{T}_{\mathbf{M}} = \{E \in \mathcal{E} : \models_{\mathbf{M}} E\}$$

Logical axioms LA are always a subset of expressions that are **tautologies** of under the semantics \mathbf{M} , i.e.

$$LA \subseteq \mathbf{T}_{\mathbf{M}}$$

Components: Rules of Inference

Rules of inference \mathcal{R}

We **assume** that a proof system contains only a **finite number** of **inference rules**

We **assume** that each rule has a **finite number** of **premisses** and **one conclusion**

We also **assume** that one can **effectively decide**, for any **inference rule**, whether a given string of expressions **form** its premisses and conclusion or **do not**, i.e. that

All rules $r \in \mathcal{R}$ are **primitively recursive**

Components: Rules of Inference

Definition

Each **rule of inference** $r \in \mathcal{R}$ is a **relation** defined in the set \mathcal{E}^m , where $m \geq 1$ with values in \mathcal{E} , i.e.

$$r \subseteq \mathcal{E}^m \times \mathcal{E}$$

Elements P_1, P_2, \dots, P_m of a tuple $(P_1, P_2, \dots, P_m, C) \in r$ are called **premisses** of the rule r and C is called its **conclusion**

All $r \in \mathcal{R}$ are **primitively recursive** relations

Components: Rules of Inference

We write the **inference rules** in a following convenient way

One premiss rule

$$(r) \frac{P_1}{C}$$

Two premisses rule

$$(r) \frac{P_1 ; P_2}{C}$$

m premisses rule

$$(r) \frac{P_1 ; P_2 ; \dots ; P_m}{C}$$

Semantic Link: Sound Rules of Inference

Given some **m** premisses rule

$$(r) \quad \frac{P_1 ; P_2 ; \dots ; P_m}{C}$$

Semantical link

Given a semantics **M** for the language \mathcal{L} and for the set of expressions \mathcal{E}

We want the **rules of inference** $r \in \mathcal{R}$ to **preserve truthfulness** i.e. to be **sound** under the semantics **M**

General Definition: Sound Rule of Inference

Definition

Given an inference rule $r \in \mathcal{R}$

$$(r) \quad \frac{P_1 ; P_2 ; \dots ; P_m}{C}$$

We say that the inference rule $r \in \mathcal{R}$ is **sound** under a semantics **M**

if and only if

all **M** - **models** of the set $\{P_1, P_2, \dots, P_m\}$ of its **premisses** are also **M** - **models** of its **conclusion C**

Propositional Definition: Sound Rule of Inference

In **propositional languages** case, the semantics **M**, and hence the **M - models** are defined in terms of the truth assignment $v : VAR \rightarrow LV$, where **LV** is the set of **logical values** for the semantics **M**

Definition

An inference rule $r \in \mathcal{R}$, such that

$$(r) \quad \frac{P_1 ; P_2 ; \dots ; P_m}{C}$$

is sound under a semantics **M**

if and only if

the condition below holds for any $v : VAR \rightarrow LV$

If $v \models_M \{P_1, P_2, \dots, P_m\}$, **then** $v \models_M C$

Example

Given a rule of inference

$$(r) \frac{(A \Rightarrow B)}{(B \Rightarrow (A \Rightarrow B))}$$

Prove that (r) is **sound** under classical semantics

Let v be any truth assignment, such that $v \models (A \Rightarrow B)$, i.e.
by definition $v^*(A \Rightarrow B) = T$

We evaluate logical value of the **conclusion** under v as follows

$$v^*(B \Rightarrow (A \Rightarrow B)) = v^*(B) \Rightarrow T = T$$

for any B and any value of $v^*(B)$

This proves that $v \models (B \Rightarrow (A \Rightarrow B))$ and hence the **soundness** of (r)

Formal Definition: Proof System

Definition

By a **proof system** we understand a quadruple

$$S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$$

where

$\mathcal{L} = \{\mathcal{A}, \mathcal{F}\}$ is a **language** of S with a set \mathcal{F} of formulas

\mathcal{E} is a set of **expressions** of S

In particular case $\mathcal{E} = \mathcal{F}$

$LA \subseteq \mathcal{E}$ is a **non- empty, finite set** of **logical axioms** of S

\mathcal{R} is a **non- empty, finite set** of **rules of inference** of S

PART 3: Formal Proofs
Simple Examples of Proof Systems

Provable Expressions

A **final product** of a **single** or **multiple** use of the **inference rules** of **S**, with **axioms** taken as a **starting point** are called **provable expressions** of the proof system **S**

A **single** use of an **inference rule** is called a **direct consequence**

A **multiple** application of rules of inference with **axioms** taken as a **starting point** is called a **proof**

Definition: Direct Consequence

Formal definitions are as follows

Direct consequence

For any rule of inference $r \in \mathcal{R}$ of the form

$$(r) \quad \frac{P_1 ; P_2 ; \dots ; P_m}{C}$$

C is called a **direct consequence** of P_1, \dots, P_m by virtue of the rule $r \in \mathcal{R}$

Definition: Formal Proof

Formal Proof of an expression $E \in \mathcal{E}$ in a proof system

$$S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$$

is a sequence

$$A_1, A_2, \dots, A_n \text{ for } n \geq 1$$

of expressions from \mathcal{E} , such that

$$A_1 \in LA, \quad A_n = E$$

and for each $1 < i \leq n$, either $A_i \in LA$ or A_i is a **direct consequence** of some of the **preceding expressions** by virtue of **one of the rules of inference**

$n \geq 1$ is the **length of the proof** A_1, A_2, \dots, A_n

Formal Proof Notation

We write

$$\vdash_S E$$

to denote that $E \in \mathcal{E}$ **has a proof** in S

When the proof system S is **fixed** we write $\vdash E$

Any $E \in \mathcal{E}$, such that $\vdash_S E$ is called a **provable expression** of S

The set of **all provable expressions** of S is denoted by \mathbf{P}_S , i.e. we put

$$\mathbf{P}_S = \{E \in \mathcal{E} : \vdash_S E\}$$

PART 4: Hypothesis, Consequence,
Soundness and Completeness

Proof from Hypothesis

While proving expressions we often use **some extra information** available, besides the axioms of the proof system. This extra information is called **hypothesis** in the proof.

Let $\Gamma \subseteq \mathcal{E}$ be a set expressions called **hypothesis**

A proof of $E \in \mathcal{E}$ **from the set of hypothesis** Γ in S is a **formal proof** in S , where the expressions from Γ are treated as **additional hypothesis added** to the set LA of the **logical axioms** of the system S

Notation: $\Gamma \vdash_S A$

We read it : A has a proof in S from the set Γ (and logical axioms LA)

Definition: Proof from Hypothesis

Definition

We say that A **has a proof** in S **from the set** Γ (and logical axioms LA) if and only if

there is a sequence A_1, \dots, A_n of expressions from \mathcal{E} , such that

$$A_1 \in LA \cup \Gamma, \quad A_n = A$$

and for each $1 < i \leq n$, either $A_i \in LA \cup \Gamma$ or A_i is a **direct consequence** of some of the **preceding expressions** by virtue of **one of the rules** of inference

We denote it as $\Gamma \vdash_S A$

Special Cases

We usually consider and use the case when the set of hypothesis is finite.

Case of $\Gamma \subseteq \mathcal{E}$ **finite set** and $\Gamma = \{B_1, B_2, \dots, B_n\}$

We use **notation**

$$B_1, B_2, \dots, B_n \vdash_S A$$

for $\{B_1, B_2, \dots, B_n\} \vdash_S A$

Case of $\Gamma = \emptyset$ is also a special one.

By the definition of a proof of A from Γ , $\emptyset \vdash A$ means that in the proof of A we use **only axioms LA** of S

We hence use **notation** $\vdash_S A$

to denote that A has a proof from **empty** Γ ; i.e. A has a proof from logical axioms only

Definition: Consequences of Γ

Definition

For any $\Gamma \subseteq \mathcal{E}$, and $A \in \mathcal{E}$,

If $\Gamma \vdash_S A$, then A is called a **consequence** of Γ in S

Definition

We denote by $\mathbf{Cn}_S(\Gamma)$ the **set of all consequences** of Γ in S , i.e. we put

$$\mathbf{Cn}_S(\Gamma) = \{A \in \mathcal{E} : \Gamma \vdash_S A\}$$

Definition: Consequence Operation

Observe that by defining a consequence of Γ in S , we define in fact a **function** which to every set $\Gamma \subseteq \mathcal{E}$ assigns a set of **all its consequences** $\mathbf{Cn}_S(\Gamma)$

We denote this function by \mathbf{Cn}_S and adopt the following

Definition

Any function

$$\mathbf{Cn}_S : 2^{\mathcal{E}} \longrightarrow 2^{\mathcal{E}}$$

such that for every $\Gamma \in 2^{\mathcal{E}}$

$$\mathbf{Cn}_S(\Gamma) = \{A \in \mathcal{E} : \Gamma \vdash_S A\}$$

is called the **consequence operation** in S

Consequence Operation: Monotonicity

Take any **consequence operation**

$$\mathbf{Cn}_S : 2^{\mathcal{E}} \longrightarrow 2^{\mathcal{E}}$$

Monotonicity Property

For any sets Γ, Δ of expressions of S ,

if $\Gamma \subseteq \Delta$ **then** $\mathbf{Cn}_S(\Gamma) \subseteq \mathbf{Cn}_S(\Delta)$

Exercise: write the proof;

it follows directly from the definition of \mathbf{Cn}_S and definition of the formal proof

Consequence Operation: Transitivity

Take any **consequence operation**

$$\mathbf{Cn}_S : 2^{\mathcal{E}} \longrightarrow 2^{\mathcal{E}}$$

Transitivity Property

For any sets $\Gamma_1, \Gamma_2, \Gamma_3$ of expressions of S ,

if $\Gamma_1 \subseteq \mathbf{Cn}_S(\Gamma_2)$ and $\Gamma_2 \subseteq \mathbf{Cn}_S(\Gamma_3)$, **then** $\Gamma_1 \subseteq \mathbf{Cn}_S(\Gamma_3)$

Exercise: write the proof;

it follows directly from the definition of \mathbf{Cn}_S and definition of the formal proof

Consequence Operation: Finiteness

Take any **consequence operation**

$$\mathbf{Cn}_S : 2^{\mathcal{E}} \rightarrow 2^{\mathcal{E}}$$

Finiteness Property

For any expression $A \in \mathcal{E}$ and any set $\Gamma \subseteq \mathcal{E}$,

$A \in \mathbf{Cn}_S(\Gamma)$ if and only if there is a **finite subset** Γ_0 of Γ such that $A \in \mathbf{Cn}_S(\Gamma_0)$

Exercise: write the proof;

it follows directly from the definition of \mathbf{Cn}_S and definition of the formal proof

Definition: Sound S

Definition

Given a proof system

$$S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$$

We say that the system **S** is **sound** under a semantics **M** iff the following conditions hold

1. $LA \subseteq T_M$
2. Each rule of inference $r \in \mathcal{R}$ is **sound**

Example

Given a proof system:

$$S = (\mathcal{L}_{\{\neg, \Rightarrow\}}, \mathcal{F}, \{(A \Rightarrow A), (A \Rightarrow (\neg A \Rightarrow B))\}, (r) \frac{(A \Rightarrow B)}{(B \Rightarrow (A \Rightarrow B))})$$

1. Prove that **S** is **sound** under **classical semantics**
2. Prove that **S** is **not sound** under **K** semantics

Example

1. Both **axioms** of **S** are basic classical **tautologies** and we have just proved that the rule of inference **(r)** is **sound**, hence **S** is **sound**

2. Axiom $(A \Rightarrow A)$ is not a **K** semantics tautology

Any truth assignment v such that $v^*(A) = \perp$ is a **counter-model** for it

This proves that **S** is **not sound** under **K** semantics

Soundness Theorem

Let \mathbf{P}_S be the set of all provable expressions of S i.e.

$$\mathbf{P}_S = \{A \in \mathcal{E} : \vdash_S A\}$$

Let \mathbf{T}_M be a set of all expressions of S that are tautologies under a semantics \mathbf{M} , i.e.

$$\mathbf{T}_M = \{A \in \mathcal{E} : \models_M A\}$$

Soundness Theorem for S and semantics \mathbf{M}

$$\mathbf{P}_S \subseteq \mathbf{T}_M$$

i.e. for any $A \in \mathcal{E}$, the following implication holds

$$\text{If } \vdash_S A, \text{ then } \models_M A.$$

Exercise: prove by Mathematical Induction over the length of a proof that if S is sound, the Soundness Theorem holds for S

Completeness Theorem

Completeness Theorem for S and semantics M

$$P_S = T_M$$

i.e. for any $A \in \mathcal{E}$, the following holds

$$\vdash_S A \quad \text{if and only if} \quad \models_M A$$

The **Completeness Theorem** consists of two parts:

Part 1: **Soundness Theorem**

$$P_S \subseteq T_M$$

Part 2: **Completeness Part** of the Completeness Theorem

$$T_M \subseteq P_S$$

Syntactic Consistency: Formal Theories

Formal theories play crucial role in **mathematics** and were historically defined for classical **predicate (first order)** logic and consequently for other non-classical logics

They are routinely called **first order theories**

We discuss them in detail in **Chapter 10** dealing formally with **classical predicate** logic

First order theories are hence based on a proof systems **S** with a **predicate** (first order) language \mathcal{L}

We sometimes consider **formal theories** based on proof systems with a **propositional** language \mathcal{L} and we call them **propositional theories**

Syntactic Consistency: Formal Theories

Given a proof system $S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$

We **build** (define) a **formal theory** based on S as follows.

1. We **select** a certain **finite** subset SA of expressions of S , **disjoint** with the logical axioms LA of S

The set SA is called a set of **specific** axioms of the **formal theory** based on S

2. We use set SA of **specific** axioms to define a language \mathcal{L}_{SA} , called a **language** of the formal theory

Here we have two cases

Syntactic Consistency: Formal Theories

c1 S is a first order proof system, i.e. \mathcal{L} of S is a **predicate** language

We **define** the language \mathcal{L}_{SA} by **restricting** the sets of **constant, functional**, and **predicate** symbols of \mathcal{L} to constant, functional, predicate symbols **appearing** in the set SA of **specific axioms**

Both languages \mathcal{L}_{SA} and \mathcal{L} **share** the same set of **propositional** connectives

c2 S is a **propositional** proof system, i.e. \mathcal{L} of S is a **propositional** language \mathcal{L}_{SA} is defined by **restricting** \mathcal{L} to connectives appearing in the set SA

Syntactic Consistency: Formal Theories

Definition

Given a proof system $S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$ and **finite** subset SA of expressions of S , **disjoint** with the logical axioms LA

The system

$$T = (\mathcal{L}, \mathcal{E}, LA, SA, \mathcal{R})$$

is called a **formal theory** based on S

The set SA is the set of **specific axioms** of T

The language \mathcal{L}_{SA} defined by **c1** or **c2** is called the language of the **theory** T

Syntactic Consistency

Definition

A theory

$$T = (\mathcal{L}, \mathcal{E}, LA, SA, \mathcal{R})$$

is **consistent** if and only if there exists an expression $E \in \mathcal{E}_{SA}$ such that $E \notin \mathbf{T}(SA)$, i.e. such that

$$SA \not\vdash_S E$$

otherwise the theory T is **inconsistent**.

Observe that the definition has **purely syntactic** meaning

Syntactic Consistency: Formal Theories

The **consistency** definition reflexes our **intuition** what proper notion of **provability** should mean

Namely, it **says** that a formal **theory** T based on a proof system S is **consistent** only when it **does not prove** all expressions (formulas in particular cases) of \mathcal{L}_{SA}

The **theory** T such that it **proves everything** stated in \mathcal{L}_{SA} obviously should be, and **is defined** as **inconsistent**

Syntactic Consistency: Formal Theories

In particular, we have the following **syntactic definition** of **consistency** and **inconsistency** for any proof system S

Definition

A proof system

$$S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$$

is **consistent** if and only if there exists $E \in \mathcal{E}$ such that $E \notin \mathbf{P}_S$, i.e. such that

$$\not\vdash_S E$$

otherwise S is **inconsistent**

Formal Theory

Given a proof system $S = (\mathcal{L}, \mathcal{E}, LA, \mathcal{R})$. Let a set $SA \subseteq \mathcal{E}$ be such that

$$SA \cap T_M = \emptyset$$

A **formal theory** with the set of **specific axioms** SA is denoted by $T(SA)$ and defined as follows

$$T(SA) = (\mathcal{L}_{SA}, \mathcal{E}, LA, SA, \mathcal{R})$$

The set of all expressions of the language \mathcal{L}_{SA} provable from the set specific axioms SA (and logical axioms LA) i.e. the set

$$T(SA) = \{A \in \mathcal{E} : SA \vdash_S A\}$$

is called the set of all **theorems** of the theory $T(SA)$

Soundness of the Theory

Soundness Theorem for a formal theory $T(SA)$ based on a proof system S says:

For any formula A of the language \mathcal{L}_{SA} of the theory $T(SA)$,
if a formula A is **provable** in the theory $T(SA)$,
then A is **true** in any **model** of the set of **specific axioms**
 SA of $T(SA)$

Syntactic Completeness of Formal Theories

The **Completeness Theorem** for the proof system **S** established equivalency of the notion of provability and tautology:

$$P_S = T_M$$

Observe the equation $P_S = T_M$ holds for a theory $T(SA)$ **only when** the set of its specific axioms $SA = \emptyset$

We nevertheless talk about **Complete/Incomplete** theories as the **final goal** of the course (and the book) is to prove the **Gödel Incompleteness Theorem** for the **Peano** formal theory of the **Arithmetic of Natural Numbers**

Complete Formal Theory

Definition

A formal theory $T(SA)$ based on a language with negation \neg is **complete** if and only if **for any** A of the language of the theory $T(SA)$ the following holds

$$A \in T(SA) \text{ or } \neg A \in T(SA)$$

Otherwise a theory $T(SA)$ is **incomplete**

The **completeness** of a theory means that we can **prove** or **disapprove** any statement **formulated within it**

It hence corresponds to the natural meaning of the notion of a **complete information**

Syntactic Consistence

Definition

A formal theory $T(SA)$ based on a language with negation \neg is **consistent** if and only if **there is no** formula A of the **language** of the theory $T(SA)$ such that

$$A \in T(SA) \text{ and } \neg A \in T(SA)$$

Otherwise $T(SA)$ is **inconsistent**

The notions of **consistency**, **inconsistency** and **completeness**, **incompleteness** describe are the most important properties of any theory

PART 5: Decidable and Syntactically Decidable Proof Systems

Decidable and Syntactically Decidable Proof Systems

A proof system S is called **decidable** when there is a **finite, mechanical method** for determining, given any expression $A \in \mathcal{E}$ whether **there is** a proof of A in S ; i.e. whether $A \in \mathbf{P}_S$

otherwise S is called **undecidable**

Observe that the above notion of decidability of the system **does not require to find a proof**

It requires only a mechanical procedure of deciding whether a **proof exists** for any expression of the system.

Example

We **prove now** that A Hilbert style proof system S for classical propositional logic presented in Chapter 9 is **decidable**

We first prove the **Completeness Theorem** for it

$$\mathbf{P}_S = \mathbf{T}_M$$

We get that for any $A \in \mathcal{E}$

$$A \notin \mathbf{P}_S \quad \text{iff} \quad A \notin \mathbf{T}_M$$

We have proved already that that the notion of classical propositional tautology, i.e. the statement $A \notin \mathbf{T}_M$ is **decidable**

We conclude: the system S is **decidable**

Syntactically Decidable Systems

A proof system S is **syntactically decidable** if it is possible to define for it a **finite, mechanical method** that **generates a proof** for any given expression A of S

otherwise the system S **is not syntactically decidable**

We call such syntactically decidable systems **automated theorem proving** systems

Syntactically Decidable Systems

All **Gentzen type** proof systems presented here are both **decidable** or **semi-decidable** and **syntactically decidable** or **syntactically semi-decidable**.

We usually call them **automated theorem proving** systems for different logics under consideration.

Resolution based proof systems are also widely known examples of the syntactically decidable, or semi-decidable systems.

Finding a Gentzen Type, or Resolution type formalization for a given logic is a **standard question** one asks about any logic being developed.

Formal Proofs

Remember that the notion of a **formal proof** in a system S is **purely syntactical** in its nature

Formal Proof carries a semantical meaning via established semantics and the **Soundness Theorem**

The **rules of inference** of a proof system define only how to **transform strings of symbols** of the language **into another string** of symbols.

The **formal proof**, by the definition says that in order to **prove** an expression A in a system S one has to construct of a **sequence** of proper transformations, **defined** by the rules of inference.

Simple System S_1

Consider a very simple proof system system S_1 with $\mathcal{E} = \mathcal{F}$

$$S_1 = (\mathcal{L}_{\{P, \Rightarrow\}}, \mathcal{F}, LA1 = \{(A \Rightarrow A)\}, (r) \frac{B}{PB}),$$

where $A, B \in \mathcal{F}$ are any formulas and

where P is some one argument connective;

we might read PA for example as "it is possible that A "

Observe that even the system S_1 has only **one axiom**, it represents an **infinite** number of formulas.

We call such axiom **axiom schema**

Simple System S_2

Consider now a system S_2

$$S_2 = (\mathcal{L}_{\{P, \Rightarrow\}}, \mathcal{F} \text{ LA2} = \{(a \Rightarrow a)\}, (r) \frac{B}{PB}),$$

where $a \in VAR$ is any variable (atomic formula) and $B \in \mathcal{F}$ is any formula

Observe that even the system S_1 has only **one axiom**, it is also an **axiom schema**

Observe that for example a formula

$$((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c)))$$

is an **axiom** of system S_1

but **is not** an axiom of the system S_2

Some Provable Formulas

We have that

$$\vdash_{S_1} ((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c)))$$

because

$$((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c))) \in \text{LA1}$$

other provable formulas are

$$\vdash_{S_1} P(a \Rightarrow a), \quad \vdash_{S_2} P(a \Rightarrow a),$$

$$\vdash_{S_1} PP(a \Rightarrow a), \quad \vdash_{S_2} PP(a \Rightarrow a)$$

Formal Proofs

Formal proofs in both systems of above formulas are identical and are as follows.

Formal proof of $P(a \Rightarrow a)$ in S_1 and S_2 is:

$A_1 = (a \Rightarrow a),$	$A_2 = P(a \Rightarrow a)$
axiom	rule application
	for $B = (a \Rightarrow a)$

Formal Proofs

Formal proof of $PP(a \Rightarrow a)$ in S_1 and S_2 is:

$A_1 = (a \Rightarrow a),$	$A_2 = P(a \Rightarrow a),$	$A_3 = PP(a \Rightarrow a)$
axiom	rule application	rule application
	for $B = (a \Rightarrow a)$	for $B = P(a \Rightarrow a)$

Proof Search

Let's **search for a proof** (if exists) of the formula **A** below in S_2

$$A = PP((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c)))$$

Observe, that if **A** had the proof, **the only last step** in this proof would be the application of the **rule** $(r) \frac{B}{PB}$ to the formula

$$P((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c)))$$

This formula, in turn, if it had the proof, **the only** last step in its proof would be the application of the **rule** r to the formula

$$((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c)))$$

The **search process stops here**

Proof Search

Observe that

$$((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c))) \notin LA2$$

what means that our **search** for the proof has **failed**;

i.e. our **found sequence** of formulas **does not constitute a proof**

Moreover, the search was, **at each step unique** what proves that the proof of **A** in **S₂** **does not exist**, i.e.

$$\not\vdash_{S_2} PP((Pa \Rightarrow (b \Rightarrow c)) \Rightarrow (Pa \Rightarrow (b \Rightarrow c)))$$

Proof Search Procedure

We easily **generalize** above example to a proof search procedure to any formula **A** of S1 or S2 as follows

Procedure SP

Step: Check the **main connective** of **A**

If **main connective** is **P**, it means that **A** was obtained by the rule **r**

Erase the main connective **P**

Repeat until no **P** as a **main connective** is left.

If the main connective is \Rightarrow check if a formula is an **axiom**

If **it is** an axiom, **stop** and **yes** we have a **proof**

If **it is not** an axiom, **stop** and **no**, proof **does not exist**

Syntactical Decidability

The **Procedure SP** is a **finite, effective, automatic** procedure of searching for a proof of formulas in both our proof systems. This proves the following.

Fact Proof systems S_1 and S_2 are **syntactically decidable**

Semantical link

Remark that we haven't defined a **semantics** for the language $\mathcal{L}_{\{\Rightarrow, P\}}$ of systems **S1, S2**

We can't talk about the **soundness** of these systems yet but we can think how to define a sound semantics for our systems.

If we want to understand statement **PA** as "A is possible" we need to define some kind of **modal** semantics.

Semantical link

All known **modal semantics extend** the classical semantics, i.e. they are **the same** as classical one on **non-modal connectives**

Hence under any possible modal semantics axioms **S1, S2** of would be a **sound axiom** under standard modal logics semantics, as they are classical tautologies.

To **assure the soundness** of both systems we must have a modal semantics **M** that makes the rule

$$(r) \frac{B}{PB}$$

sound under the modal semantics **M**

General Question 1

General Q1: Are all proof systems decidable?

Answer Q1: No, not all proof systems are decidable

The most "natural" and historically first developed proof system for **classical predicate logic** is **not decidable**

General Question 2

General Q2 Can we give an **example** of a logic and its complete proof system which **is not decidable**, but the logic does have another complete, **syntactically decidable** proof system?

Answer Q2: Hilbert style proof system for classical propositional logic presented in chapter 5 is **complete** and **decidable** but **is not syntactically decidable**

We present in chapter 6 **some complete proof systems** for classical propositional logic that **are syntactically decidable**