

Discrete Mathematics

Chapter 5 Problem 16

Chapter 4 Problem 14

Chapter 5, Problem No 16

- Evaluate the sum

$$\sum_k \begin{pmatrix} 2a \\ a+k \end{pmatrix} \begin{pmatrix} 2b \\ b+k \end{pmatrix} \begin{pmatrix} 2c \\ c+k \end{pmatrix} (-1)^k$$

Continued...

The binomial coefficient $\binom{n}{k}$ can be

expressed in terms of factorials as follows:

$$\binom{n}{k} = n! / (k! * (n - k)!)$$

Continued...

Lets try to express each of the terms in the problem in factorials :

$$\binom{2a}{a+k} = \frac{(2a)!}{(2a-(a+k))! * (a+k)!}$$
$$= \frac{(2a)!}{(a-k)! * (a+k)!}$$

Continued...

Similarly,

$$\binom{2b}{b+k} = \frac{(2b)!}{(b-k)! * (b+k)!}$$

$$\binom{2c}{c+k} = \frac{(2c)!}{(c-k)! * (c+k)!}$$

Continued...

Therefore,

$$\sum_k \binom{2a}{a+k} \binom{2b}{b+k} \binom{2c}{c+k} (-1)^k$$

$$= \frac{\sum_k (2a)! (2b)! (2c)! (-1)^k}{k (a-k)! (a+k)! (b-k)! (b+k)! (c-k)! (c+k)!}$$

Continued...

Multiplying numerator and denominator by
 $(a+b)! * (b+c)! * (c+a)!$

We will therefore have,

$$\frac{(2a)! (2b)! (2c)!}{(a+b)! (b+c)! (c+a)!} * \sum_k \frac{(a+b)! (b+c)! (c+a)! (-1)^k}{(a-k)! (a+k)! (b-k)! (b+k)! (c-k)! (c+k)!}$$

Constant

Lets try to get a known form for this.

Continued...

Considering :

$$\sum_k \frac{(a+b)! (b+c)! (c+a)!}{(a-k)! (a+k)! (b-k)! (b+k)! (c-k)! (c+k)!}$$

$$= \sum_k \frac{(a+b)! (b+c)! (c+a)!}{(a+k)! (b-k)! (b+k)! (c-k)! (c+k)! (a-k)!}$$

(Just interchanging the order of the terms in the denominator)

We know that,

$$\frac{(a+b)!}{(a+k)! (b-k)!} = \binom{a+b}{a+k}$$

Continued...

Similarly,

$$\frac{(b+c)!}{(b+k)! (c-k)!} = \binom{b+c}{b+k}$$

$$\frac{(c+a)!}{(c+k)! (a-k)!} = \binom{c+a}{c+k}$$

Continued...

Therefore,

$$\sum_k \frac{(a+b)! (b+c)! (c+a)! * (-1)^k}{(a-k)! (a+k)! (b-k)! (b+k)! (c-k)! (c+k)!}$$

$$= \sum_k \binom{a+b}{a+k} \binom{b+c}{b+k} \binom{c+a}{c+k} * (-1)^k$$

which is a known form.

Using the equation given in Textbook Page No . 171,Eq. 5-29.

We have,

$$\sum_k \binom{a+b}{a+k} \binom{b+c}{b+k} \binom{c+a}{c+k} * (-1)^k = \frac{(a + b + c)!}{a! b! c!}$$

Solution

Thus, the solution for the problem becomes :

$$\frac{(2a)! (2b)! (2c)!}{(a+b)! (b+c)! (c+a)!} * \frac{(a+b+c)!}{a! b! c!}$$

Chapter 4, Problem No 14

Does every prime occur as a factor of some Euclid number e_n ?

Continued...

Euclid Number :

Definition :

Euclid numbers are integers of the

form $E_n = p_n\# + 1$,

where $p_n\#$ is the primorial of p_n which is the n th prime.

Continued...

They are named after the ancient Greek mathematician Euclid, who used them in his original proof that there are an infinite number of prime numbers.

Primorial :

For $n \geq 2$, the **primorial** ($n\#$) is the product of all prime numbers less than or equal to n . For example, $7\# = 210$ is a primorial which is the product of the first four primes multiplied together ($2 \cdot 3 \cdot 5 \cdot 7$).

Continued...

The simplest argument could be that to show that there is a prime number which is never the factor of any Euclid number.

If we consider any Euclid number,

$p_n\#$ is always a multiple of 2.

And Euclid number is 1 added to $p_n\#$.

Continued..

Every Euclid number is of the form
$$= (2 * k) + 1$$

where “k” is product of prime numbers $\leq n$
excluding 2.

So, it is very clear that there exists no
Euclid number which is divisible by 2.

Answer

Hence, the answer is :

Every prime cannot occur as a factor of some Euclid number e_n .



THANK YOU