# cse581 COMPUTER SCIENCE FOUNDAMENTALS: THEORY

Professor Anita Wasilewska

▲□▶ ▲□▶ ▲ □▶ ▲ □▶ ▲ □ ● のへで

Lecture 1

# DISCRETE MATHEMATICS BASICS

#### **Discrete Mathematics Basics**

PART 0: Basic sets of Numbers, Peano Arithmetic

- PART 1: Sets and Operations on Sets
- PART 2: Relations and Functions
- PART 3: Special types of Binary Relations
- PART 4: Finite and Infinite Sets
- PART 5: Some Fundamental Proof Techniques

## Theory of Computation BASICS

- PART 6: Closures and Algorithms
- PART 7: Alphabets and languages
- PART 8: Finite Representation of Languages

(日)

**Discrete Mathematics Basics** 

PART 0: Basic sets of Numbers, Peano Arithmetic

Basic Sets of Numbers

Natural numbers N, Integers Z, Positive Integers  $Z^+$ , Positive Natural numbers  $N^+$ , Prime Numbers P, Rational Numbers Q, and Real numbers R Natural Numbers N

 $N = \{0, 1, 2, 3, \ldots, \ldots\}$ 

Integers Z and Positive Integers  $Z^+$ 

 $Z = \{ \ldots, -3, -2, -1, 0, 1, 2, 3, \ldots, \ldots \}$ 

 $Z^+ = \{ 1, 2, 3, \ldots, \ldots \}$ 

Positive Integers  $Z^+$  are also called Positive Natural numbers  $N^+$  and we denote

$$N^+ = \{ 1, 2, 3, \ldots, \ldots \}$$

・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・
 ・

### **Prime Numbers**

A positive integer  $p \in Z^+$  is called prime if it has only two divisors, namely 1 and p By convention, 1 **is not** prime

Prime Numbers P

 $P = \{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, \dots, \dots\}$ 

### **Rational and Real Numbers**

Rational numbers Q

$$Q = \{\frac{p}{q}: p, q \in Z \text{ and } q \neq 0\}$$

Real numbers R

The **first rigorous definition** of the set **R** of real numbers was published by **Cantor** in 1871

**Cantor's definition** (as established today in modern terminology)

The set R Is the **quotient set** of the set of Cauchy sequences of **rational numbers**, with two sequences considered **equivalent** if their difference converges to zero

Cantor also showed In 1874, that the set of all real numbers is uncountably infinite, but the set of all algebraic numbers is countably infinite

#### **Real Numbers**

The **other first rigorous definition** of **R** established today was given by **Richard Dedekind** at the same time and independent from **Cantor** in terms what we call now **Dedekind cuts** 

The concept of the Dedekind cuts developed for it became on of the very important concepts for modern mathematics

The set of **R** is often called "The Reals" - after the name "real numbers" first used by a French philosopher, scientist, and mathematician **Rene Descartes** (1596 -1650), also known as **Renatus Cartesius** 

#### Irrational and Algebraic Numbers

Of course we have that  $N \subset Q \subset R$ Real numbers that are not Rational are called **Irrational** numbers, i.e. we put IR = R - Q**Algebraic number** is a number that is a **root** of a non-zero polynomial P(x) in one variable equation P(x) = 0 with **integer** (or, equivalently **rational**) coefficients

#### All rational numbers are algebraic

Let  $x \in Q$ , by the definition  $x = \frac{a}{b}$  for any integers  $a, b \neq 0$  is the root of a non-zero polynomial equation namely bx - a = 0

### Encyclopedia Britannica

Here is what is published the Encyclopedia Britannica

**Real number** in mathematics, is a quantity that can be expressed as an infinite decimal expansion

The real numbers include the positive and negative integers and the fractions made from those integers (or rational numbers) and**also** the irrational numbers

Natural Numbers in Encyclopedia Britannica

Here is what is published the Encyclopedia Britannica

**Natural numbers:** called the counting numbers or natural numbers (1, 2, 3, ). For an empty set, no object is present, and the **count yields** the number 0, which, appended to the counting numbers, produces what are known as the **whole numbers** Hence the Modern Mathematics definition is

 $N = \{0, 2, 3, 4, \dots \}$ 

 $N = Z^+ \cup \{0\} = N^+ \cup \{0\} =$  whole numbers

Next to geometry, the **theory of natural numbers** is the most intuitive and intuitively known of all branches of mathematics

This is why the first attempts to **formalize mathematics** begin with arithmetic of natural numbers.

(日)

The first attempt of axiomatic formalization was given by Dedekind in **1879** and by Peano in **1889** 

The Peano formalization became known as **Peano Postulates** and can be written as follows.

## Peano Postulates (1889)

p1 0 is a natural number

**p2** If *n* is a natural number, there is another number which we denote by n'We call the number n' a **successor** of *n* and the intuitive meaning of n' is n + 1

**p3**  $0 \neq n'$ , for any natural number **n** 

**p4** If n' = m', then n = m, for any natural numbers n, m

p5 If W is is a property that may or may not hold for natural numbers, and
if (i) 0 has the property W and
(ii) whenever a natural number n has the property W, then n' has the property W,
then all natural numbers have the property W

The postulate **p5** is called Principle of Induction

The **Peano Postulates** together with certain amount of set theory are sufficient to develop **not only** theory of natural numbers, **but also** theory of rational and even real numbers

But **Peano Postulates** can't act as a fully formal theory as they include **intuitive** notions like "property" and

"has a property" . A **formal theory** of natural numbers based on the Peano

Postulates is referred in literature as **Peano Arithmetic**, or simply **PA** 

We present, in Chapter 11 of the book **B2** a formalization by Mendelson (1973) It is included and worked out in the smallest **details** in his book *Intoduction to Mathematical Logic* (1987) **Discrete Mathematics Basics** 

(ロト (個) (E) (E) (E) (9)

PART 1: Sets and Operations on Sets

### Sets

Set A set is a collection of objects

**Elements** The objects comprising a set are are called its elements or members

 $a \in A$  denotes that a is an **element** of a set A

 $a \notin A$  denotes that a is not an **element** of A

Empty Set is a set without elements

Empty Set is denoted by Ø

#### Sets

Sets can be defined by listing their elements;

Example

The set

$$A = \{a, \emptyset, \{a, \emptyset\}\}$$

has 3 elements:

 $a \in A$ ,  $\emptyset \in A$ ,  $\{a, \emptyset\} \in A$ 

### Sets

Sets can be defined by referring to other sets and to properties P(x) that elements may or may not have

We write it as

 $B = \{x \in A : P(x)\}$ 

#### Example

Let N be a set of natural numbers

 $B = \{n \in N : n < 0\} = \emptyset$ 

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

#### Set Inclusion

 $A \subseteq B$  if and only if  $\forall a (a \in A \Rightarrow a \in B)$ is a **true** statement

Set Equality A = B if and only if  $A \subseteq B$  and  $B \subseteq A$ 

**Proper Subset**  $A \subset B$  if and only if  $A \subseteq B$  and  $A \neq B$ 

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

#### **Subset Notations**

- $A \subseteq B$  for a subset (might be improper)  $A \subset B$  for a proper subset
- **Power Set** Set of all subsets of a given set

 $\mathcal{P}(A) = \{B : B \subseteq A\}$ 

**Other Notation** 

$$2^{A} = \{B : B \subseteq A\}$$

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

#### Union

 $A \cup B = \{x : x \in A \text{ or } x \in B\}$ 

We write:

 $x \in A \cup B$  if and only if  $x \in A \cup x \in B$ 

# Intersection $A \cap B = \{x : x \in A \text{ and } x \in B\}$ We write: $x \in A \cap B$ if and only if $x \in A \cap x \in B$

◆□▶ ◆□▶ ◆三▶ ◆三▶ ・三 のへで

#### **Relative Complement**

 $x \in (A - B)$  if and only if  $x \in A$  and  $x \notin B$ We write:

$$A-B=\{x: x\in A \cap x \notin B\}$$

**Complement** is defined only for  $A \subseteq U$ , where *U* is called an **universe** 

$$-A = U - A$$

▲□▶ ▲圖▶ ▲匡▶ ▲匡▶ ― 匡 - のへで

We write for  $x \in U$ ,

 $x \in -A$  if and only if  $x \notin A$ 

Algebra of sets consists of properties of sets that are true for all sets involved

We use **tautologies** of propositional logic to prove **basic** properties of the algebra of sets

We then use the basic properties to **prove** more elaborated properties of sets

(日)

It is possible to form intersections and unions of **more** then two, or even a finite number o **sets** 

Let  $\mathcal{F}$  denote is any **collection** of sets

We write  $\bigcup \mathcal{F}$  for the set whose elements are the elements of all of the sets in  $\mathcal{F}$ 

Example Let

 $\mathcal{F} = \{\{a\}, \{\emptyset\}, \{a, \emptyset, b\}\}$ 

We get

$$\bigcup \mathcal{F} = \{a, \ \emptyset, \ b\}$$

Observe that given

$$\mathcal{F} = \{\{a\}, \{\emptyset\}, \{a, \emptyset, b\}\} = \{A_1, A_2, A_3\}$$

we have that

 $\{a\} \cup \{\emptyset\} \cup \{a, \emptyset, b\} = A_1 \cup A_2 \cup A_3 = \{a, \emptyset, b\} = \left( \begin{array}{c} \int \mathcal{F} \\ \int \mathcal{F} \\ \end{array} \right)$ 

Hence we have that for any element x,

 $x \in \bigcup \mathcal{F}$  if and only if there exists i, such that  $x \in A_i$ 

We **define** formally **Generalized Union** of any family  $\mathcal{F}$  of sets is

 $\int \mathcal{F} = \{x : \text{ exists a set } S \in \mathcal{F} \text{ such that } x \in S\}$ 

We write it also as

$$x \in \bigcup \mathcal{F}$$
 if and only if  $\exists_{S \in \mathcal{F}} x \in S$ 

▲□▶▲□▶▲□▶▲□▶ ■ のへで

Generalized Intersection of any family  $\ensuremath{\mathcal{F}}$  of sets is

$$\bigcap \mathcal{F} = \{ x : \forall_{S \in \mathcal{F}} x \in S \}$$

We write

$$x \in \bigcap \mathcal{F}$$
 if and only if  $\forall_{S \in \mathcal{F}} x \in S$ 

▲□▶▲圖▶▲≣▶▲≣▶ ≣ のQ@

#### **Ordered Pair**

Given two sets A, B we denote by

# (a, b)

an **ordered pair**, where  $a \in A$  and  $b \in B$ We call a a **first** coordinate of (a, b)and b its **second** coordinate We define

(a,b) = (c,d) if and only if a = c and b = d

▲□▶▲□▶▲□▶▲□▶ □ のQ@

## **Cartesian Product**

Given two sets A and B, the set

 $A \times B = \{(a, b) : a \in A \text{ and } b \in B\}$ 

is called a **Cartesian product** (cross product) of sets *A*, *B* We write

 $(a, b) \in A \times B$  if and only if  $a \in A$  and  $b \in B$ 

▲□▶▲□▶▲□▶▲□▶ □ のQ@

**Discrete Mathematics Basics** 

PART 2: Relations and Functions

## **Binary Relations**

## **Binary Relation**

Any set R such that  $R \subseteq A \times A$ is called a **binary relation** defined in a set A

# **Domain, Range** of R Given a binary relation $R \subseteq A \times A$ , the set

 $D_R = \{a \in A : (a, b) \in R\}$ 

is called a domain of the relation R

The set

$$V_R = \{b \in A : (a, b) \in R\}$$

is called a range (set of values) of the relation R

#### n- ary Relations

#### Ordered tuple

Given sets  $A_1, ..., A_n$ , an element  $(a_1, a_2, ..., a_n)$  such that  $a_i \in A_i$  for i = 1, 2, ..., n is called an **ordered tuple** 

**Cartesian Product** of sets  $A_1, A_n$  is a set

 $A_1 \times A_2 \times ... \times A_n = \{(a_1, a_2, ..., a_n) : a_i \in A_i, i = 1, 2, ..., n\}$ 

**n-ary Relation** on sets  $A_1, \ldots, A_n$  is any subset of  $A_1 \times A_2 \times \ldots \times A_n$ , i.e. the set

 $R \subseteq A_1 \times A_2 \times \ldots \times A_n$ 

## **Binary Relations**

## **Binary Relation**

Any set R such that  $R \subseteq A \times B$ is called a **binary relation** defined in a sets A and B

# **Domain, Range** of R Given a binary relation $R \subseteq A \times B$ , the set

 $D_R = \{a \in A : (a, b) \in R\}$ 

is called a domain of the relation R

The set

```
V_R = \{b \in B : (a, b) \in R\}
```

is called a range (set of values) of the relation R

## Function as Relation

## Definition

A binary relation  $R \subseteq A \times B$  on sets A, B is a **function** from A to B

if and only if the following condition holds

 $\forall_{a\in A} \exists! _{b\in B} (a,b) \in R$ 

where  $\exists !_{b \in B}$  means there is **exactly one**  $b \in B$ 

Because the condition says that for any  $a \in A$  we have **exactly one**  $b \in B$ , we write

R(a) = b for  $(a, b) \in R$ 

Function as Relation

Given a binary relation

 $R\subseteq A\times B$ 

that is a **function** 

The set *A* is called a **domain** of the function *R* and we write:

$$R: A \longrightarrow B$$

▲ロト ▲ 同 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()

to denote that the relation R is a function and say that

*R* maps the set A into the set B
## **Function notation**

We denote relations that are functions by letters  $f, g, h, \dots$  and write

 $f: A \longrightarrow B$ 

▲□▶▲□▶▲□▶▲□▶ □ のQ@

say that the function f maps the set A into the set B

Domain, Codomain

Let  $f: A \longrightarrow B$ ,

the set A is called a **domain** of f,

and the set B is called a codomain of f

## Range

Given a function  $f: A \longrightarrow B$ 

The set

 $R_f = \{b \in B : b = f(a) \text{ and } a \in A\}$ 

is called a **range** of the function f

By definition, the **range** of f is a subset of its **codomain** B We write  $R_f = \{b \in B : \exists_{a \in A} b = f(a)\}$ 

The set

$$f = \{(a, b) \in A \times B : b = f(a)\}$$

▲ロト ▲ 同 ト ▲ ヨ ト ▲ ヨ ト ・ ヨ ・ の Q ()

is called a graph of the function f

Function "onto"

The function  $f: A \longrightarrow B$  is an **onto** function if and only if the following condition holds

 $\forall_{b\in B} \exists_{a\in A} f(a) = b$ 

We denote it by

 $f: A \xrightarrow{onto} B$ 

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

Function "one-to-one"

The function  $f: A \longrightarrow B$ is called a **one- to -one** function and denoted by

 $f: A \xrightarrow{1-1} B$ 

if and only if the following condition holds

 $\forall_{x,y\in A} (x\neq y \Rightarrow f(x)\neq f(y))$ 

▲□▶▲□▶▲□▶▲□▶ □ のQ@

A function  $f: A \longrightarrow B$  is **not one- to -one** function if and only if the following condition holds

 $\exists_{x,y\in A}(x\neq y\cap f(x)=f(y))$ 

If a function f is **1-1** and **onto** we denote it as

 $f: A \xrightarrow{1-1,onto} B$ 

## **Composition of functions**

Let f and g be two functions such that

 $f: A \longrightarrow B$  and  $g: B \longrightarrow C$ 

We define a new function

 $h: A \longrightarrow C$ 

called a **composition** of functions f and g as follows: for any  $x \in A$  we put

h(x) = g(f(x))

▲□▶▲□▶▲□▶▲□▶ □ のQ@

## **Composition notation**

Given function f and g such that

 $f: A \longrightarrow B$  and  $g: B \longrightarrow C$ 

We **denote** the **composition** of f and g by  $(f \circ g)$  in order to stress that the function

 $f: A \longrightarrow \mathbf{B}$ 

"goes first" followed by the function

 $g: \mathbf{B} \longrightarrow C$ 

with a shared set B between them

We write now the **definition** of composition of functions **f** and **g** using the **composition notation** (name for the composition function)  $(f \circ g)$  as follows The composition  $(f \circ g)$  is a **new** function

 $(f \circ g) : A \longrightarrow C$ 

such that for any  $x \in A$  we put

 $(f \circ g)(x) = g(f(x))$ 

▲□▶▲□▶▲□▶▲□▶ □ のQ@

There is also other notation (name) for the **composition** of f and g that uses the symbol  $(g \circ f)$ , i.e. we put

 $(g \circ f)(x) = g(f(x))$  for all  $x \in A$ 

This notation was invented to help calculus students to remember the formula g(f(x)) defining the composition of functions f and g

▲□▶▲□▶▲□▶▲□▶ □ のQ@

## **Inverse function**

Let  $f: A \longrightarrow B$  and  $g: B \longrightarrow A$ 

*g* is called an **inverse** function to *f* if and only if the following condition holds

# $\forall_{a\in A}(f\circ g)(a)=g(f(a))=a$

▲□▶▲□▶▲□▶▲□▶ □ のQ@

If g is an **inverse** function to f we denote by  $g = f^{-1}$ 

## **Identity function**

A function  $I: A \longrightarrow A$  is called an **identity** on A if and only if the following condition holds

 $\forall_{a\in A} l(a) = a$ 

#### **Inverse and Identity**

Let  $f : A \longrightarrow B$  and let  $f^{-1} : B \longrightarrow A$ be an **inverse** to f, then the following hold

 $(f \circ f^{-1})(a) = f^{-1}(f(a)) = I(a) = a$ , for all  $a \in A$ 

 $(f^{-1} \circ f(b)) = f(f^{-1}(b) = l(b) = b$ , for all  $b \in B$ 

▲□▶▲圖▶▲≣▶▲≣▶ = ● ● ●

Functions: Image and Inverse Image

#### Image

Given a function  $f: X \longrightarrow Y$  and a set  $A \subseteq X$ The set

$$f[A] = \{y \in Y : \exists x \ (x \in A \cap y = f(x))\}$$

is called an **image** of the set  $A \subseteq X$  under the function f We write

 $y \in f[A]$  if and only if there is  $x \in A$  and y = f(x)

Other symbols used to denote the image are

$$f^{\rightarrow}(A)$$
 or  $f(A)$ 

◆□▶ ◆□▶ ◆ □▶ ◆ □▶ ○ □ ○ ○ ○ ○

Functions: Image and Inverse Image

## Inverse Image

Given a function  $f: X \longrightarrow Y$  and a set  $B \subseteq Y$ The set

$$f^{-1}[B] = \{x \in X : f(x) \in B\}$$

is called an **inverse image** of the set  $B \subseteq Y$  under the function f

We write

$$x \in f^{-1}[B]$$
 if and only if  $f(x) \in B$ 

Other symbol used to denote the inverse image are

$$f^{-1}(B)$$
 or  $f^{\leftarrow}(B)$ 

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

#### Sequences

#### Definition

A **sequence** of elements of a set A is any **function** from the set of natural numbers N into the set A, i.e. any function

 $f: N \longrightarrow A$ 

Any  $f(n) = a_n$  is called **n-th term** of the sequence f Notations  $\{a_n\}_{n \in \mathbb{N}}, \{a_n\}_{n \ge 0}$ 

 $a_0, a_1, a_2, \ldots, \ldots$ 

## Sequences

We often consider sequences  $\{a_n\}_{n\geq 1}$  and adopt **Definition** 

A **sequence** of elements of a set A is any **function** from the set of positive natural numbers  $N^+$  or from the set positive Integers  $Z^+$  into the set A, i.e. any function

$$f: \mathbb{N}^+ \longrightarrow \mathbb{A}$$
 or  $f: \mathbb{Z}^+ \longrightarrow \mathbb{A}$ 

Any  $f(n) = a_n$  is called **n-th term** of the sequence f Notations  $\{a_n\}_{n \in \mathbb{Z}^+}, \{a_n\}_{n \in \mathbb{N}^+}, \{a_n\}_{n \ge 1}$ 

 $a_1, a_2, a_3, \ldots, \ldots$ 

## Sequences Example

#### Example

We define a sequence f of real numbers R as follows

 $f: N \longrightarrow R$ 

such that

$$f(n)=n+\sqrt{n}$$

We also use a shorthand notation for the function f and write it as

 $a_n = n + \sqrt{n}$ 

▲□▶▲□▶▲□▶▲□▶ ■ のへで

#### Sequences Example

We often write the function  $f = \{a_n\}$  in an even shorter and **informal** form as

 $a_0 = 0$ ,  $a_1 = 1 + 1 = 2$ ,  $a_2 = 2 + \sqrt{2}$ .....

or even as

0, 2, 2 +  $\sqrt{2}$ , 3 +  $\sqrt{3}$ , ..... n +  $\sqrt{n}$ .....

▲□▶▲□▶▲□▶▲□▶ □ のQ@

## Observations

## **Observation 1**

By definition, **sequence** of elements of any set is always infinite (countably infinite) because the domain of the **sequence** function f is a set N of **natural numbers** 

## **Observation 2**

We can enumerate elements of a sequence by any infinite subset of  $\ensuremath{\mathsf{N}}$ 

We often take a set  $N^+ = N - \{0\}$  as a **sequence** domain (enumeration) and "start" with n = 1, i.e. write

$$a_1, a_2, a_3, \dots, a_n, \dots$$

#### Observations

#### **Observation 3**

We can choose as a set of indexes of a **sequence** any countably infinite set T, i. e, **not only** the set N of natural numbers We often choose  $T = N - \{0\} = N^+$ , i.e we consider **sequences** that "start" with n = 1In this case we write sequences as

 $a_1, a_2, a_3, \dots, a_n, \dots$ 

## Finite Sequences

## **Finite Sequence**

Given a finite set  $K = \{1, 2, ..., n\}$ , for  $n \in N$  and any set A

Any function

 $f: \{1, 2, ..., n\} \longrightarrow A$ 

is called a **finite sequence** of elements of the set A of the **length** n

## Case n=0

In this case the function f is an empty set and we call it an

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

#### empty sequence

We denote the empty sequence by e

Other common notation is  $\lambda$ 

## Example

#### Example

Consider a sequence given by a formula

$$a_n=\frac{n}{(n-2)(n-5)}$$

The domain of the function  $f(n) = a_n$  is the set  $N - \{2, 5\}$ and the **sequence** f is a function

 $f: N-\{2,5\} \rightarrow R$ 

The first elements of the sequence f are

 $a_0 = f(0), \ a_1 = f(1), \ a_3 = f(3), \ a_4 = f(4), \ a_6 = f(6), \dots$ 

## Families of Sets

## Family of sets

Any collection of sets is called a **family of sets** We denote the family of sets by  $\mathcal{F}$ Sequence of sets Any function

 $f: N \longrightarrow \mathcal{F} \text{ or } f: N^+ \longrightarrow \mathcal{F}$ 

is a **sequence of sets**, i..e a sequence where all its elements are sets

We use capital letters to denote sets and write the **sequence** of sets as  $\{A_n\}_{n \in \mathbb{N}}$ ,  $\{A_n\}_{n \in \mathbb{N}^+}$ ,  $\{A_n\}_{n \geq 1}$ 

#### **Generalized Union**

#### **Generalized Union**

Given a sequence  $\{A_n\}_{n \in \mathbb{N}}$  of sets

We define that Generalized Union of the sequence of sets as

$$\bigcup_{n\in N} A_n = \{x : \exists_{n\in N} x \in A_n\}$$

We write

$$x \in \bigcup_{n \in \mathbb{N}} A_n$$
 if and only if  $\exists_{n \in \mathbb{N}} x \in A_n$ 

**Generalized Intersection** 

**Generalized Intersection** 

Given a sequence  $\{A_n\}_{n \in \mathbb{N}}$  of sets We define that **Generalized Intersection** of the sequence of sets as

$$\bigcap_{n\in\mathbb{N}}A_n=\{x: \forall_{n\in\mathbb{N}} x\in A_n\}$$

We write

$$x \in \bigcap_{n \in N} A_n$$
 if and only if  $\forall_{n \in N} x \in A_n$ 

◆□▶ ◆□▶ ◆ □▶ ◆ □▶ ○ □ ○ ○ ○ ○

# Indexed Family of Sets

## **Indexed Family of Sets**

Given  $\mathcal{F}$  be a family of sets Let  $T \neq \emptyset$  be any non empty set

## Any function

 $f: T \longrightarrow \mathcal{F}$ 

is called an indexed family of sets with the set of indexes T We write it

# $\{\mathbf{A}_t\}_{t\in T}$

#### Notice

Any sequence of sets is an indexed family of sets for T = N

▲□▶▲□▶▲□▶▲□▶ □ のQ@

Short Review

# Some Simple Questions and Answers

・ロト・日本・日本・日本・日本・日本

Simple Short Questions

Here are some short **Yes**/ **No** questions Answer them and write a short **justification** of your answer

- **Q1**  $2^{\{1,2\}} \cap \{1,2\} \neq \emptyset$
- **Q2**  $\{\{a, b\}\} \in 2^{\{a, b, \{a, b\}\}}$
- **Q3**  $\emptyset \in 2^{\{a,b,\{a,b\}\}}$
- **Q4** Any function f from  $A \neq \emptyset$  onto A, has property

 $f(a) \neq a$  for certain  $a \in A$ 

## Simple Short Questions

**Q5** Let  $f: N \longrightarrow \mathcal{P}(N)$  be given by a formula:  $f(n) = \{m \in N : m < n^2\}$ 

then  $\emptyset \in f[\{0, 1, 2\}]$ 

# **Q6** Some relations $R \subseteq A \times B$

are functions that map the set A into the set B

▲□▶ ▲□▶ ▲ 三▶ ▲ 三▶ - 三 - のへぐ

Q1  $2^{\{1,2\}} \cap \{1,2\} \neq \emptyset$ NO because

 $2^{\{1,2\}} = \{\emptyset, \{1\}, \{2\}, \{1,2\}\} \cap \{1,2\} = \emptyset$ 

Q2  $\{\{a, b\}\} \in 2^{\{a, b, \{a, b\}\}}$ YES because have that  $\{a, b\} \subseteq \{a, b, \{a, b\}\}$  and hence  $\{\{a, b\}\} \in 2^{\{a, b, \{a, b\}\}}$ 

by definition of the set of all subsets of a given set

Q2  $\{\{a, b\}\} \in 2^{\{a, b, \{a, b\}\}}$ YES other solution We list all subsets of the set  $\{a, b, \{a, b\}\}$ , i.e. all elements of the set

2<sup>{a,b,{a,b}}</sup>

We start as follows

```
\{\emptyset, \{a\}, \{b\}, \{\{a, b\}\}, \ldots, \ldots\}
```

and observe that we can **stop** listing because we reached the set  $\{\{a, b\}\}\$ This proves that  $\{\{a, b\}\} \in 2^{\{a, b, \{a, b\}\}}$ 

- **Q3**  $\emptyset \in 2^{\{a,b,\{a,b\}\}}$
- **YES** because for any set A, we have that  $\emptyset \subseteq A$
- **Q4** Any function f from  $A \neq \emptyset$  onto A has a property

 $f(a) \neq a$  for certain  $a \in A$ 

#### NO

Take a function such that f(a) = a for all  $a \in A$ Obviously f is "onto" and and there is no  $a \in A$ for which  $f(a) \neq a$ 

◆□ ▶ ◆□ ▶ ◆ 三 ▶ ◆ 三 ● ● ● ●

**Q5** Let  $f: N \to \mathcal{P}(N)$  be given by formula:  $f(n) = \{m \in N : m < n^2\}$ , then  $\emptyset \in f[\{0, 1, 2\}]$  **YES** We evaluate  $f(0) = \{m \in N : m < 0\} = \emptyset$   $f(1) = \{m \in N : m < 1\} = \{0\}$   $f(2) = \{m \in N : m < 2^2\} = \{0, 1, 2, 3\}$ and so by definition of f[A] get that  $f[\{0, 1, 2\}] = \{\emptyset, \{0\}, \{0, 1, 2, 3\}\}$  and hence  $\emptyset \in f[\{0, 1, 2\}]$ 

**Q6** Some  $R \subseteq A \times B$  are functions that map A into B **YES**: Functions are special type of relations

## Simple Short Questions

**Q7**  $\{(1,2), (a,1)\}$  is a binary relation on  $\{1,2\}$ 

**Q8** For any binary relation  $R \subseteq A \times A$ , the **inverse** relation  $R^{-1}$  **exists** 

**Q9** For any **binary relation**  $R \subseteq A \times A$  that is a function, the **inverse function**  $R^{-1}$  exists

#### Simple Short Questions

**Q10** Let  $A = \{a, \{a\}, \emptyset\}$  and  $B = \{\emptyset, \{\emptyset\}, \emptyset\}$ there is a function  $f : A \longrightarrow_{onto}^{1-1} B$ 

**Q11** Let  $f: A \rightarrow B$  and  $g: B \rightarrow onto A$ , then the compositions  $(g \circ f)$  and  $(f \circ g)$  exist

**Q12** The function  $f: N \longrightarrow \mathcal{P}(R)$  given by the formula:

$$f(n) = \{x \in R : x > \frac{ln(n^3 + 1)}{\sqrt{n + 6}}\}$$

▲□▶▲□▶▲□▶▲□▶ □ のQ@

is a **sequence** 

- **Q7**  $\{(1,2), (a,1)\}$  is a binary relation on  $\{1,2\}$
- **NO** because  $(a, 1) \notin \{1, 2\} \times \{1, 2\}$
- **Q8** For any binary relation  $R \subseteq A \times A$ , the inverse relation  $R^{-1}$  exists

**YES** By definition, the **inverse relation** to  $R \subseteq A \times A$  is the set

$$R^{-1} = \{(b, a) : (a, b) \in R\}$$

▲□▶▲□▶▲□▶▲□▶ □ のQ@

and it is a well defined relation in the set A

**Q9** For any **binary relation**  $R \subseteq A \times A$  that is a function, the **inverse function**  $R^{-1}$  exists

**NO** R must be also a 1 - 1 and *onto* function

**Q10** Let  $A = \{a, \{a\}, \emptyset\}$  and  $B = \{\emptyset, \{\emptyset\}, \emptyset\}$ there is a function  $f : A \longrightarrow_{onto}^{1-1} B$ **NO** The set A has **3** elements and the set

 $\boldsymbol{B} = \{\emptyset, \{\emptyset\}, \emptyset\} = \{\emptyset, \{\emptyset\}\}$ 

has 2 elements and an onto function does not exists
Answers to Short Questions

**Q11** Let  $f: A \longrightarrow B$  and  $g: B \longrightarrow {}^{onto} A$ , then the compositions  $(g \circ f)$  and  $(f \circ g)$  exist

**YES** The composition  $(f \circ g)$  exists because the functions  $f: A \rightarrow B$  and  $g: B \rightarrow Onto A$  share the same set B

- ロト・日本・日本・日本・日本・日本

The composition  $(g \circ f)$  exists because the functions  $g: B \longrightarrow {}^{onto} A$  and  $f: A \longrightarrow B$  share the same set A

The information "onto" is irrelevant

## Answers to Short Questions

**Q12** The function  $f: N \longrightarrow \mathcal{P}(R)$  given by the formula:

$$f(n) = \{x \in R : x > \frac{ln(n^3 + 1)}{\sqrt{n + 6}}\}$$

## is a **sequence**

**YES** It is a sequence as the **domain** of the function f is the set N of natural numbers and the formula for f(n) assigns to each natural number n a certain **subset** of R, i.e. an **element** of  $\mathcal{P}(R)$