

The Logic of Quantified Statements

CSE 215, Foundations of Computer Science

Stony Brook University

<http://www.cs.stonybrook.edu/~cse215>

The Logic of Quantified Statements

All men are mortal.

Socrates is a man.

∴ Socrates is mortal.

- *Propositional calculus*: analysis of ordinary compound statements
- *Predicate calculus*: symbolic analysis of predicates and **quantified statements** ($\forall x, \exists x$)
 - P is a *predicate symbol*
 - P stands for “is a student at SBU”
 - $P(x)$ stands for “ x is a student at SBU”
 - x is a *predicate variable*

Predicates and Quantified Statements

- A *predicate* is a sentence that contains a finite number of variables and becomes a *statement* when specific values are substituted for the variables.
- The *domain* of a predicate variable is the set of all values that may be substituted in place of the variable.
- Example:

$P(x)$ is the predicate “ $x^2 > x$ ”, x has as a domain the set \mathbf{R} of all real numbers

$P(2): 2^2 > 2.$ *True.*

$P(1/2): (1/2)^2 > 1/2.$ *False.*

Truth Set of a Predicate

- If $P(x)$ is a predicate and x has domain D , the truth set of $P(x)$, $\{x \in D \mid P(x)\}$, is the set of all elements of D that make $P(x)$ true when they are substituted for x .

- Example:

$Q(n)$ is the predicate for “ n is a factor of 8.”

if the domain of n is the set \mathbf{Z} of all integers

The truth set is $\{1, 2, 4, 8, -1, -2, -4, -8\}$

The Universal Quantifier: \forall

- Quantifiers are words that refer to quantities (“some” or “all”) and tell for how many elements a given predicate is true.
- **Universal quantifier:** \forall is “for all”
- Example:

“All human beings are mortal”

\forall human beings x , x is mortal.

- If H is the set of all human beings

$\forall x \in H$, x is mortal

Universal statements

- A **universal statement** is a statement of the form “ $\forall x \in D, Q(x)$ ” where $Q(x)$ is a predicate and D is the domain of x .
 - $\forall x \in D, Q(x)$ is true if, and only if, $Q(x)$ is true for every x in D
 - $\forall x \in D, Q(x)$ is false if, and only if, $Q(x)$ is false for at least one x in D (the value for x is a *counterexample*)
- Example:

$$\forall x \in D, x^2 \geq x \quad \text{where } D = \{1, 2, 3, 4, 5\}$$

$$1^2 \geq 1, \quad 2^2 \geq 2, \quad 3^2 \geq 3, \quad 4^2 \geq 4, \quad 5^2 \geq 5$$

- Hence “ $\forall x \in D, x^2 \geq x$ ” is true.

The Existential Quantifier: \exists

- **Existential quantifier:** \exists is “there exists” or “for some”

- Example:

- “There is a student in CSE 215”

\exists a person p such that p is a student in CSE 215

$\exists p \in P$ (a person p), such that, p is a student in CSE 215

where P is the set of all people

The Existential Quantifier: \exists

- An **existential statement** is a statement of the form “ $\exists x \in D$ such that $Q(x)$ ” where $Q(x)$ is a predicate and D the domain of x
 - $\exists x \in D$ s.t. $Q(x)$ is true if, and only if, $Q(x)$ is true for at least one x in D
 - $\exists x \in D$ s.t. $Q(x)$ is false if, and only if, $Q(x)$ is false for all x in D
- Example:
 - $\exists m \in \mathbb{Z}$ such that $m^2 = m$
 $1^2 = 1$ True
- Notation: I will use s.t. for "such that" to be concise

The Relation among \forall , \exists , \wedge , and \vee

- $D = \{x_1, x_2, \dots, x_n\}$ and

$$\forall x \in D, Q(x) \equiv Q(x_1) \wedge Q(x_2) \wedge \dots \wedge Q(x_n)$$

- $D = \{x_1, x_2, \dots, x_n\}$ and

$$\exists x \in D \text{ such that } Q(x) \equiv Q(x_1) \vee Q(x_2) \vee \dots \vee Q(x_n)$$

Universal Conditional Statements

- **Universal conditional statement:**


$\forall x, \text{ if } P(x) \text{ then } Q(x)$

- Example:

If a real number is greater than 2 then its square is greater than 4.

$\forall x \in \mathbf{R}, \text{ if } x > 2 \text{ then } x^2 > 4$

Vacuous Truth of Universal Statements



All the balls in the bowl are blue

True

$\forall x$ in D , if $P(x)$ then $Q(x)$ is *vacuously true* or *true by default* if, and only if, $P(x)$ is false for every x in D

Equivalent Forms of Universal and Existential Statements

- $\forall x \in U, \text{ if } P(x) \text{ then } Q(x)$ can be rewritten in the form $\forall x \in D, Q(x)$ by narrowing U to be the domain D consisting of all values of the variable x that make $P(x)$ true.
 - Example: $\forall x, \text{ if } x \text{ is a square then } x \text{ is a rectangle}$
 $\forall \text{ squares } x, x \text{ is a rectangle.}$
- $\exists x \text{ such that } P(x) \text{ and } Q(x)$ can be rewritten in the form $\exists x \in D \text{ such that } Q(x)$ where D consists of all values of the variable x that make $P(x)$ true

Implicit Quantification

- $P(x) \Rightarrow Q(x)$ means that every element in the truth set of $P(x)$ is in the truth set of $Q(x)$, or, equivalently, $\forall x, P(x) \rightarrow Q(x)$
- $P(x) \Leftrightarrow Q(x)$ means that $P(x)$ and $Q(x)$ have identical truth sets, or, equivalently, $\forall x, P(x) \boxed{\leftrightarrow} Q(x)$.

Negations of Quantified Statements

- Negation of a Universal Statement:

The negation of a statement of the form $\forall x \in D, Q(x)$ is logically equivalent to a statement of the form

$\exists x \in D, \sim Q(x)$, that is:

$$\sim(\forall x \in D, Q(x)) \equiv \exists x \in D, \sim Q(x)$$

- Example:
 - “All mathematicians wear glasses”
 - Its negation is: “There is at least one mathematician who does not wear glasses”
 - IMPORTANT: Its negation is NOT “No mathematicians wear glasses”!!!!

Negations of Quantified Statements

- Negation of an Existential Statement

The negation of a statement of the form $\exists x \in D, Q(x)$ is logically equivalent to a statement of the form

$\forall x \in D, \sim Q(x)$, that is:

$$\sim(\exists x \in D, Q(x)) \equiv \forall x \in D, \sim Q(x)$$

- Example:

- “Some snowflakes are the same.”

- Its negation is: “No snowflakes are the same” which means: “All snowflakes are different.”

Negations of Quantified Statements

- More Examples:
 - $\sim(\forall \text{ primes } p, p \text{ is odd}) \equiv \exists \text{ a prime } p \text{ such that } p \text{ is **not** odd}$
 - $\sim(\exists \text{ a triangle } T \text{ such that the sum of the angles of } T \text{ equals } 200^\circ) \equiv \forall \text{ triangles } T, \text{ the sum of the angles of } T \text{ **does not** equal } 200^\circ$
 - $\sim(\forall \text{ politicians } x, x \text{ is **not** honest}) \equiv \exists \text{ a politician } x \text{ such that } x \text{ is honest (**by double negation**)}$
 - $\sim(\forall \text{ computer programs } p, p \text{ is finite}) \equiv \exists \text{ a computer program } p \text{ that is not finite}$
 - $\sim(\exists \text{ a computer hacker } c, c \text{ is over } 40) \equiv \forall \text{ computer hacker } c, c \text{ is } 40 \text{ or under}$
 - $\sim(\exists \text{ an integer } n \text{ between } 1 \text{ and } 37 \text{ such that } 1,357 \text{ is divisible by } n) \equiv \forall \text{ integers } n \text{ between } 1 \text{ and } 37, 1,357 \text{ is not divisible by } n$

Negations of Quantified Statements

- $\sim(\forall x, P(x) \rightarrow Q(x)) \equiv \exists x \text{ such that } P(x) \wedge \sim Q(x)$

Proof:

$$\sim(\forall x, P(x) \rightarrow Q(x)) \equiv \exists x \text{ such that } \sim(P(x) \rightarrow Q(x))$$

$$\sim(P(x) \rightarrow Q(x)) \equiv \sim(\sim P(x) \vee Q(x)) \equiv \sim\sim P(x) \wedge \sim Q(x) \equiv P(x) \wedge \sim Q(x)$$

- Examples:
 - $\sim(\forall \text{ people } p, \text{ if } p \text{ is blond then } p \text{ has blue eyes}) \equiv$
 $\exists \text{ a person } p \text{ such that } p \text{ is blond and } p \text{ does not have blue eyes}$
 - $\sim(\text{If a computer program has more than 100,000 lines, then it contains a bug})$
 $\equiv \text{There is at least one computer program that has more than 100,000 lines}$
 $\text{and does not contain a bug}$

Variants of Universal Conditional Statements

- Universal conditional statement: $\forall x \in D$, if $P(x)$ then $Q(x)$
- **Contrapositive:** $\forall x \in D$, if $\sim Q(x)$ then $\sim P(x)$

$\forall x \in D$, if $P(x)$ then $Q(x) \equiv \forall x \in D$, if $\sim Q(x)$ then $\sim P(x)$

Proof: for any x in D by the logical equivalence between statement and its contrapositive

- **Converse:** $\forall x \in D$, if $Q(x)$ then $P(x)$.
- **Inverse:** $\forall x \in D$, if $\sim P(x)$ then $\sim Q(x)$.
- Example:

$\forall x \in \mathbb{R}$, if $x > 2$ then $x^2 > 4$

Contrapositive: $\forall x \in \mathbb{R}$, if $x^2 \leq 4$ then $x \leq 2$

Converse: $\forall x \in \mathbb{R}$, if $x^2 > 4$ then $x > 2$

Inverse: $\forall x \in \mathbb{R}$, if $x \leq 2$ then $x^2 \leq 4$

Necessary and Sufficient Conditions

- Necessary condition:

“ $\forall x, r(x)$ is a **necessary condition** for $s(x)$ ” means

“ $\forall x, \text{if } \sim r(x) \text{ then } \sim s(x)$ ” \equiv “ $\forall x, \text{if } s(x) \text{ then } r(x)$ ” (*)

(*)(by contrapositive and double negation)

- Sufficient condition:

“ $\forall x, r(x)$ is a **sufficient condition** for $s(x)$ ” means

“ $\forall x, \text{if } r(x) \text{ then } s(x)$ ”

Necessary and Sufficient Conditions

- Examples:

- Squareness is a **sufficient condition** for rectangularity;

Formal statement: $\forall x$, if x is a square, then x is a rectangle

- Being at least 35 years old is a **necessary condition** for being President of the United States

\forall people x , if x is younger than 35, then x cannot be President of the United States \equiv

\forall people x , if x is President of the United States then x is at least 35 years old (by contrapositive)

Only If

- Only If:

“ $\forall x, r(x)$ **only if** $s(x)$ ” means

“ $\forall x, \text{if } \sim s(x) \text{ then } \sim r(x)$ ” \equiv “ $\forall x, \text{if } r(x) \text{ then } s(x)$.”

- Example:

A product of two numbers is 0 **only if** one of the numbers is 0.

If neither of two numbers is 0, then the product of the numbers is not 0 \equiv

If a product of two numbers is 0, then one of the numbers is 0 (by contrapositive)

Statements with Multiple Quantifiers

- Example:

“There is a person supervising every detail of the production process”

- What is the meaning?

“There is one single person who supervises all the details of the production process”?

OR

“For any particular production detail, there is a person who supervises that detail, but there might be different supervisors for different details”?

NATURAL LANGUAGE IS AMBIGUOUS

LOGIC IS CLEAR


Statements with Multiple Quantifiers

- Quantifiers are performed in the order in which the quantifiers occur.
- Example:

$\forall x$ in set D, $\exists y$ in set E such that x and y satisfy property $P(x, y)$


Tarski's World

- Blocks of various shapes, and colors located on a grid



- $\forall t, \text{Triangle}(t) \rightarrow \text{Blue}(t)$ TRUE
- $\forall x, \text{Blue}(x) \rightarrow \text{Triangle}(x).$ FALSE
- $\exists z$ such that $\text{Square}(z) \wedge \text{Gray}(z).$ FALSE

Statements with Multiple Quantifiers in Tarski's World




$\forall \exists$

- For all triangles x , there is a square y such that x and y have the same color
TRUE

Given $x =$	choose $y =$	and check that y is the same color as x .
d	e	yes ✓
f or i	h or g	yes ✓

Statements with Multiple Quantifiers in Tarski's World



$\exists \forall$

- There is a triangle x such that for all circles y , x is to the right of y


TRUE

Choose $x =$	Then, given $y =$	check that x is to the right of y .
d or i	a	yes ✓
	b	yes ✓
	c	yes ✓

Interpreting Statements with Two Different Quantifiers


- $\forall x$ in D , $\exists y$ in E such that $P(x, y)$
 - for whatever element x in D you must find an element y in E that “works” for that particular x
- $\exists x$ in D such that $\forall y$ in E , $P(x, y)$
 - find one particular x in D that will “work” no matter what y in E anyone might choose

Interpreting Statements with Two Different Quantifiers



- \exists an item I such that \forall students S , S chose I . TRUE
- \exists a student S such that \forall stations Z , \exists an item I in Z such that S chose I TRUE
- \forall students S and \forall stations Z , \exists an item I in Z such that S chose I . FALSE

Statements with Multiple Quantifiers in Tarski's World




$\forall \exists$

- For all triangles x , there is a square y such that x and y have the same color
TRUE

Given $x =$	choose $y =$	and check that y is the same color as x .
d	e	yes ✓
f or i	h or g	yes ✓

Statements with Multiple Quantifiers in Tarski's World



$\exists \forall$

- There is a triangle x such that for all circles y , x is to the right of y .

TRUE

Choose $x =$	Then, given $y =$	check that x is to the right of y .
d or i	a	yes ✓
	b	yes ✓
	c	yes ✓

Negations of Multiply-Quantified Statements

- Apply negation to quantified statements from left to right:

$$\sim(\forall x \text{ in } D, \exists y \text{ in } E \text{ such that } P(x, y))$$

$$\equiv \exists x \text{ in } D \text{ such that } \sim(\exists y \text{ in } E \text{ such that } P(x, y))$$


$$\equiv \exists x \text{ in } D \text{ such that } \forall y \text{ in } E, \sim P(x, y).$$

$$\sim(\exists x \text{ in } D \text{ such that } \forall y \text{ in } E, P(x, y))$$

$$\equiv \forall x \text{ in } D, \sim(\forall y \text{ in } E, P(x, y))$$

$$\equiv \forall x \text{ in } D, \exists y \text{ in } E \text{ such that } \sim P(x, y).$$

Negating Statements in Tarski's World



- For all squares x , there is a circle y such that x and y have the same color


Negation:

\exists a square x such that $\sim(\exists$ a circle y such that x and y have the same color)

$\equiv \exists$ a square x such that \forall circles y , x and y do not have the same color

TRUE: Square e is black and no circle is black.

Negating Statements in Tarski's World



- There is a triangle x such that for all squares y , x is to the right of y


Negation:

\forall triangles $x, \sim (\forall$ squares y, x is to the right of $y)$

$\equiv \forall$ triangles x, \exists a square y such that x is not to the right of y

TRUE

Quantifier Order in Tarski's World



- For every square x there is a triangle y such that x and y have different colors


TRUE

- There exists a triangle y such that for every square x , x and y have different colors

FALSE

Formalizing Statements in Tarski's World

- Triangle(x) means “x is a triangle”
- Circle(x) means “x is a circle”
- Square(x) means “x is a square”
- Blue(x) means “x is blue”
- Gray(x) means “x is gray”
- Black(x) means “x is black”
- Above(x, y) means “x is above y” (strictly above)
- RightOf(x, y) means “x is to the right of y” (strictly)
- SameColorAs(x, y) means “x has the same color as y”
- $x = y$ denotes the predicate “x is equal to y”



Formalizing Statements in Tarski's World

- For all circles x , x is above f


$$\forall x(\text{Circle}(x) \rightarrow \text{Above}(x, f))$$

- Negation:

$$\sim(\forall x(\text{Circle}(x) \rightarrow \text{Above}(x, f)))$$

$$\equiv \exists x \sim (\text{Circle}(x) \rightarrow \text{Above}(x, f))$$

$$\equiv \exists x(\text{Circle}(x) \wedge \sim \text{Above}(x, f))$$



Formalizing Statements in Tarski's World

- There is a square x such that x is black


$$\exists x(\text{Square}(x) \wedge \text{Black}(x))$$

- Negation:

$$\sim(\exists x(\text{Square}(x) \wedge \text{Black}(x)))$$

$$\equiv \forall x \sim (\text{Square}(x) \wedge \text{Black}(x))$$











$$\equiv \forall x(\sim \text{Square}(x) \vee \sim \text{Black}(x))$$



Formalizing Statements in Tarski's World

- For all circles x , there is a square y such that x and y have the same color

$$\forall x(\text{Circle}(x) \rightarrow \exists y(\text{Square}(y) \wedge \text{SameColor}(x, y)))$$


- Negation:

$$\begin{aligned} & \sim(\forall x(\text{Circle}(x) \rightarrow \exists y(\text{Square}(y) \wedge \text{SameColor}(x, y)))) \\ & \equiv \exists x \sim (\text{Circle}(x) \rightarrow \exists y(\text{Square}(y) \wedge \text{SameColor}(x, y))) \\ & \equiv \exists x(\text{Circle}(x) \wedge \sim(\exists y(\text{Square}(y) \wedge \text{SameColor}(x, y)))) \\ & \equiv \exists x(\text{Circle}(x) \wedge \forall y(\sim(\text{Square}(y) \wedge \text{SameColor}(x, y)))) \\ & \equiv \exists x(\text{Circle}(x) \wedge \forall y(\sim\text{Square}(y) \vee \sim\text{SameColor}(x, y))) \end{aligned}$$

Formalizing Statements in Tarski's World

- There is a square x such that for all triangles y , x is to right of y

$$\exists x(\text{Square}(x) \wedge \forall y(\text{Triangle}(y) \rightarrow \text{RightOf}(x, y)))$$



- Negation:

$$\sim(\exists x(\text{Square}(x) \wedge \forall y(\text{Triangle}(y) \rightarrow \text{RightOf}(x, y))))$$

$$\equiv \forall x \sim (\text{Square}(x) \wedge \forall y(\text{Triangle}(y) \rightarrow \text{RightOf}(x, y)))$$

$$\equiv \forall x(\sim \text{Square}(x) \vee \sim(\forall y(\text{Triangle}(y) \rightarrow \text{RightOf}(x, y))))$$

$$\equiv \forall x(\sim \text{Square}(x) \vee \exists y(\sim(\text{Triangle}(y) \rightarrow \text{RightOf}(x, y))))$$

$$\equiv \forall x(\sim \text{Square}(x) \vee \exists y(\text{Triangle}(y) \wedge \sim \text{RightOf}(x, y)))$$

Arguments with Quantified Statements

- ***Universal instantiation***: if some property is true of everything in a set, then it is true of any particular thing in the set.
- Example:
 - All men are mortal.
 - \therefore If Socrates is a man, then Socrates is mortal.

Universal Modus Ponens

Formal Version

$\forall x$, if $P(x)$ then $Q(x)$.

$P(a)$ for a particular a .

$\therefore Q(a)$.

- Example:

$\forall x$, if $E(x)$ then $S(x)$.

$E(k)$, for a particular k .

$\therefore S(k)$.

Informal Version

If x makes $P(x)$ true, then x makes

$Q(x)$ true.

a makes $P(x)$ true.

$\therefore a$ makes $Q(x)$ true.

If an integer is even, then its square is even.

k is a particular integer that is even.

$\therefore k^2$ is even.

Universal Modus Tollens

Formal Version

$\forall x$, if $P(x)$ then $Q(x)$.
 $\sim Q(a)$, for a particular a .
 $\therefore \sim P(a)$.

- Example:

$\forall x$, if $H(x)$ then $M(x)$
 $\sim M(Z)$
 $\therefore \sim H(Z)$.

Informal Version

If x makes $P(x)$ true, then x makes
 $Q(x)$ true.
 a does not make $Q(x)$ true.
 $\therefore a$ does not make $P(x)$ true.

All human beings are mortal.
Zeus is not mortal.
 \therefore Zeus is not human.

Validity of Arguments with Quantified Statements


- An argument form is **valid**, if and only if, for any particular predicates substituted for the predicate symbols in the premises **if the resulting premise statements are all true, then the conclusion is also true**
- Using Diagrams to Test for Validity

Using Diagrams to Test for Validity

All human beings are mortal.

Zeus is not mortal.

∴ Zeus is not a human being.




Using Diagrams to Show Invalidity

All human beings are mortal.

Felix is mortal.

∴ Felix is a human being.




Using Diagrams to Test for Validity

- Universal modus tollens Example:

No polynomial functions have horizontal asymptotes.

This function has a horizontal asymptote.

∴ This function is not a polynomial function



Universal Transitivity

Formal Version

$\forall x P(x) \rightarrow Q(x).$

$\forall x Q(x) \rightarrow R(x).$

$\therefore \forall x P(x) \rightarrow R(x).$

• Example from Tarski's World:

$\forall x$, if x is a triangle, then x is blue.

$\forall x$, if x is blue, then x is to the right of all the squares.

$\therefore \forall x$, if x is a triangle, then x is to the right of all the squares

Informal Version

Any x that makes $P(x)$ true makes $Q(x)$ true.

Any x that makes $Q(x)$ true makes $R(x)$ true.

\therefore Any x that makes $P(x)$ true makes $R(x)$ true.

Converse Error (in Quantified Form)

Formal Version

$\forall x, \text{ if } P(x) \text{ then } Q(x).$

$Q(a)$ for a particular a .

$\therefore P(a).$

invalid conclusion

Informal Version

If x makes $P(x)$ true, then x makes
 $Q(x)$ true.

a makes $Q(x)$ true.

$\therefore a$ makes $P(x)$ true.

Inverse Error (in Quantified Form)

Formal Version

$\forall x$, if $P(x)$ then $Q(x)$.

$\sim P(a)$, for a particular a .

$\therefore \sim Q(a)$.

invalid conclusion

Informal Version

If x makes $P(x)$ true, then x makes $Q(x)$ true.

a does not make $P(x)$ true.

$\therefore a$ does not make $Q(x)$ true.